

Deformed $\mathcal{W}_N$ algebras from elliptic $sl(N)$ algebras

J. Avan

LPTHE, CNRS-URA 280, Universités Paris VI/VII, France

L. Frappat, M. Rossi, P. Sorba

*Laboratoire de Physique Théorique ENSLAPP, CNRS-URA 1436,
École Normale Supérieure de Lyon and Université de Savoie, France*

Abstract

We extend to the $sl(N)$ case the results that we previously obtained on the construction of $\mathcal{W}_{q,p}$ algebras from the elliptic algebra $\mathcal{A}_{q,p}(\widehat{sl}(2)_c)$. The elliptic algebra $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$ at the critical level $c = -N$ has an extended center containing trace-like operators $t(z)$. Families of Poisson structures indexed by $N(N-1)/2$ integers, defining q -deformations of the $\mathcal{W}_N$ algebra, are constructed. The operators $t(z)$ also close an exchange algebra when $(-p^{\frac{1}{2}})^{NM} = q^{-c-N}$ for $M \in \mathbb{Z}$. It becomes Abelian when in addition $p = q^{Nh}$ where h is a non-zero integer. The Poisson structures obtained in these classical limits contain different q -deformed $\mathcal{W}_N$ algebras depending on the parity of h , characterizing the exchange structures at $p \neq q^{Nh}$ as new $\mathcal{W}_{q,p}(sl(N))$ algebras.

Résumé

Nous étendons au cas $sl(N)$ les résultats que nous avons obtenus précédemment concernant la construction des algèbres $\mathcal{W}_{q,p}$ à partir de l'algèbre elliptique $\mathcal{A}_{q,p}(\widehat{sl}(2)_c)$. L'algèbre elliptique $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$ au niveau critique $c = -N$ possède un centre étendu contenant des opérateurs de trace $t(z)$. On construit sur ce centre des familles de structures de Poisson indicées par $N(N-1)/2$ entiers, définissant des q -déformations de l'algèbre $\mathcal{W}_N$. Les opérateurs $t(z)$ engendrent une algèbre d'échange lorsque $(-p^{\frac{1}{2}})^{NM} = q^{-c-N}$ où $M \in \mathbb{Z}$. Cette algèbre devient abélienne si de plus $p = q^{Nh}$ avec h entier non nul. Les structures de Poisson obtenues dans ces limites classiques contiennent différentes algèbres $\mathcal{W}_N$ q -déformées dépendant de la parité de l'entier h , caractérisant les structures d'échange à $p \neq q^{Nh}$ comme de nouvelles algèbres $\mathcal{W}_{q,p}(sl(N))$.

1 Introduction

The elliptic algebra $\mathcal{A}_{q,p}(\widehat{sl}(2)_c)$ was introduced in [1] and further studied in [2, 3, 4] where several trigonometric limits were derived and shown to be relevant as symmetries for the XXZ model, whereas $\mathcal{A}_{q,p}(\widehat{sl}(2)_c)$ was proposed as basic symmetry algebra for the XYZ model [5, 6]. More recently another, possibly related elliptic algebra $\mathcal{U}_{q,p}(\widehat{sl}(2)_c)$ was introduced [7] in connection with the k -fusion RSOS model [8, 9]. A dynamical version of the algebra $\mathcal{A}_{q,p}(\widehat{sl}(2)_c)$, making use of the dynamical “eight-vertex” elliptic R -matrix [10, 11], was also recently proposed in [12]. The domain was unified recently [13, 14] where both $\mathcal{A}_{q,p}(\widehat{sl}(2)_c)$ algebra (and its extension to $sl(N)$) and Felder’s elliptic dynamical algebra [11, 15] were shown to be obtained from application of universal twisting operators (see also [16]) to the quantum algebra $\mathcal{U}_q(\widehat{sl}(2)_c)$. As a consequence the construction [1] was validated and extended to the $sl(N)$ case.

The q -Virasoro algebra was introduced in [17] as an extension to the Ruijsenaar–Schneider model of the Virasoro algebra arising in the collective theory formulation of the Calogero–Moser model [18, 19, 20]. It arose simultaneously as quantization of a classical quadratic Poisson structure on the center at $c = -2$ of the quantum affine algebra $\mathcal{U}_q(sl(2)_c)$ [21]. It was shown [22, 23] to be the symmetry algebra of the restricted ABF model, itself connected as previously said to the elliptic $\mathcal{U}_{q,p}(\widehat{sl}(2)_c)$ algebra. On the other hand, the limit of q -Virasoro current coincides [22, 2] with the field obtained by concatenation of vertex operators of the degeneracy limit of $\mathcal{A}_{q,p}(\widehat{sl}(2)_c)$, hinting indeed to a deep connection between the two structures. This identification was recently extended to the full q -deformed Virasoro case [24].

A direct connection was indeed established in two recent papers [25, 26]: we derived exchange algebras and limit Poisson structures from the original $\mathcal{A}_{q,p}(\widehat{sl}(2)_c)$ algebra, by establishing the existence of a center at $p^m = q^{c+2}$ for any integer m . The Poisson structures all boil down to the q -Virasoro (classical) algebra defined in [17, 21]. The exchange algebras are however quite distinct from the quantization of q -Virasoro constructed in [17, 27]. Their eventual central (and linear ?) extensions also exhibit a much richer structure than was originally derived in [27] and are currently under investigation [28].

It is known that q -Virasoro structures admit extensions to the $sl(N)$ case as classical and quantum q - $\mathcal{W}_N$ algebras [21, 27, 29]. These have been the object of many investigations recently [30, 31]. In particular, they were also shown to be in general symmetry algebras of RSOS models [32, 23]. It is a natural question in view of our previous results to investigate whether classical and quantized q - $\mathcal{W}_N$ algebras arise as structures embedded in the generalized elliptic algebra $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$.

The plan of the paper is as follows. We first recall in Sect. 2 useful properties of elliptic functions and elliptic $sl(N)$ R -matrix, and introduce the definition of the quantum elliptic algebra $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$ [14]. We then prove in Sect. 3 the existence of an extended center at $c = -N$ and compute the Poisson structures on this center, using the notion of sector-depending mode Poisson bracket already introduced in [25]. These structures are identified as q -deformed $\mathcal{W}_N$ algebras. They are obtained from analytic continuations of a classical algebra identical to the initial version of q - $\mathcal{W}_N$ given in [21]. In Sect. 4 we show the existence of closed (quadratic) exchange algebras whenever $(-p^{\frac{1}{2}})^{NM} = q^{-c-N}$ for any integer $M \in \mathbb{Z}$. These algebras differ from the quantum $\mathcal{W}_{q,p}(\widehat{sl}(N))$ structures introduced in [27]. They admit a classical limit (commuting algebras) at $p = q^{Nh}$ with $h \in \mathbb{Z} \setminus \{0\}$. In Sect. 5 we compute the related Poisson structures. They include for h even the Poisson structures in [21]. The exchange algebras therefore realize new quantizations of these Poisson structures. When h is odd, by contrast, this classical limit takes a form different from the initial q - $\mathcal{W}_N$ structures. This emphasizes the key role of the initial 3-parameter structure $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$ in allowing for an

intermediate quantum q -deformed $\mathcal{W}_N$ algebra. Finally in Sect. 6, we compute the mode expansion of the quantum exchange algebra structures. As in the classical case, a “sector-type” labeling is needed due to the singularity behaviour of the structure function viewed as an analytic continuation and therefore exhibiting different formal series expansions corresponding to different convergent series expansions in distinct domains of the complex plane. We give an explicit example of this treatment applied to the spin one field for the sake of simplicity, and we describe the essential features of the extension of our computation to higher spin fields.

2 Notations and basic definitions

2.1 Definition of the N -elliptic R -matrix

The N -elliptic R -matrix in $\text{End}(\mathbb{C}^N) \otimes \text{End}(\mathbb{C}^N)$, associated to the $\mathbb{Z}_N$ -vertex model, is defined as follows [33, 34]:

$$\tilde{R}(z, q, p) = z^{2/N-2} \frac{1}{\kappa(z^2)} \frac{\vartheta \left[\begin{smallmatrix} \frac{1}{2} \\ \frac{1}{2} \end{smallmatrix} \right] (\zeta, \tau)}{\vartheta \left[\begin{smallmatrix} \frac{1}{2} \\ \frac{1}{2} \end{smallmatrix} \right] (\xi + \zeta, \tau)} \sum_{(\alpha_1, \alpha_2) \in \mathbb{Z}_N \times \mathbb{Z}_N} W_{(\alpha_1, \alpha_2)}(\xi, \zeta, \tau) I_{(\alpha_1, \alpha_2)} \otimes I_{(\alpha_1, \alpha_2)}^{-1}, \quad (2.1)$$

where the variables z, q, p are related to the variables ξ, ζ, τ by

$$z = e^{i\pi\xi}, \quad q = e^{i\pi\zeta}, \quad p = e^{2i\pi\tau}. \quad (2.2)$$

The Jacobi theta functions with rational characteristics $\vartheta \left[\begin{smallmatrix} \gamma_1 \\ \gamma_2 \end{smallmatrix} \right] (\xi, \tau)$ are defined in Appendix A.

The normalization factor is chosen as follows:

$$\frac{1}{\kappa(z^2)} = \frac{(q^{2N} z^{-2}; p, q^{2N})_\infty (q^2 z^2; p, q^{2N})_\infty (p z^{-2}; p, q^{2N})_\infty (p q^{2N-2} z^2; p, q^{2N})_\infty}{(q^{2N} z^2; p, q^{2N})_\infty (q^2 z^{-2}; p, q^{2N})_\infty (p z^2; p, q^{2N})_\infty (p q^{2N-2} z^{-2}; p, q^{2N})_\infty}. \quad (2.3)$$

The functions $W_{(\alpha_1, \alpha_2)}$ are given by

$$W_{(\alpha_1, \alpha_2)}(\xi, \zeta, \tau) = \frac{\vartheta \left[\begin{smallmatrix} \frac{1}{2} + \alpha_1/N \\ \frac{1}{2} + \alpha_2/N \end{smallmatrix} \right] (\xi + \zeta/N, \tau)}{N \vartheta \left[\begin{smallmatrix} \frac{1}{2} + \alpha_1/N \\ \frac{1}{2} + \alpha_2/N \end{smallmatrix} \right] (\zeta/N, \tau)}. \quad (2.4)$$

The matrices $I_{(\alpha_1, \alpha_2)}$ are defined as follows:

$$I_{(\alpha_1, \alpha_2)} = g^{\alpha_2} h^{\alpha_1}, \quad (2.5)$$

where the $N \times N$ matrices g and h are given by $g_{ij} = \omega^i \delta_{ij}$ and $h_{ij} = \delta_{i+1, j}$, the addition of indices being understood modulo N .

Let us set

$$S(\xi, \zeta, \tau) = \sum_{(\alpha_1, \alpha_2) \in \mathbb{Z}_N \times \mathbb{Z}_N} W_{(\alpha_1, \alpha_2)}(\xi, \zeta, \tau) I_{(\alpha_1, \alpha_2)} \otimes I_{(\alpha_1, \alpha_2)}^{-1}. \quad (2.6)$$

The matrix S is $\mathbb{Z}_N$ -symmetric, that is

$$S_{c+s, d+s}^{a+s, b+s} = S_{c, d}^{a, b} \quad (2.7)$$

for any indices $a, b, c, d, s \in \mathbb{Z}_N$ (the addition of indices being understood modulo N) and the non-vanishing elements of the matrix S are $S_{c, a+b}^{a, c+b}$. One finds explicitly:

$$S_{c, a+b}^{a, c+b} = \sum_{s \in \mathbb{Z}_N} W_{(a-c, s)} \omega^{-bs}. \quad (2.8)$$

The matrix S being $\mathbb{Z}_N$ -symmetric, it is sufficient to examine the terms $S^{ab} \equiv S_{a, b}^{0, a+b}$. One finds:

$$S^{ab}(\xi, \zeta, \tau) = \frac{\vartheta \left[\begin{smallmatrix} (b-a)/N + \frac{1}{2} \\ \frac{1}{2} \end{smallmatrix} \right] (\xi + \zeta, N\tau)}{\vartheta \left[\begin{smallmatrix} -a/N + \frac{1}{2} \\ \frac{1}{2} \end{smallmatrix} \right] (\zeta, N\tau)} \frac{\prod_{k=0, k \neq b}^{N-1} \vartheta \left[\begin{smallmatrix} k/N + \frac{1}{2} \\ \frac{1}{2} \end{smallmatrix} \right] (\xi, N\tau)}{\prod_{k=1}^{N-1} \vartheta \left[\begin{smallmatrix} k/N + \frac{1}{2} \\ \frac{1}{2} \end{smallmatrix} \right] (0, N\tau)}. \quad (2.9)$$

It satisfies the following shift properties:

$$S^{ab}(\xi, \zeta + \lambda\tau, \tau) = \exp(-2i\pi\lambda\xi/N) S^{a-\lambda, b}(\xi, \zeta, \tau), \quad (2.10a)$$

$$S^{ab}(\xi + \lambda\tau, \zeta, \tau) = \exp(-i\pi\lambda^2\tau - 2i\pi\lambda(\xi + \zeta/N + \frac{1}{2})) S^{a, b+\lambda}(\xi, \zeta, \tau). \quad (2.10b)$$

Using the following “gluing” formula:

$$\prod_{k=0}^{N-1} \vartheta \left[\begin{smallmatrix} k/N + \frac{1}{2} \\ \frac{1}{2} \end{smallmatrix} \right] (\xi, N\tau) = p^{\frac{1}{24}(N^2-1)} \frac{\prod_{k=0}^{N-1} (p^{N-k}; p^N)_\infty}{(p^N; p^N)_\infty^N} \vartheta \left[\begin{smallmatrix} \frac{1}{2} \\ \frac{1}{2} \end{smallmatrix} \right] (\xi, \tau), \quad (2.11)$$

and inserting eq. (2.9) into eqs. (2.6) and (2.1), one then finds the following expression in terms of the Jacobi Θ functions for the $\tilde{R}^{ab}$ elements of the matrix (2.1):

$$\tilde{R}^{ab}(z, q, p) = \frac{1}{\kappa(z^2)} p^{-\frac{ab}{N}} q^{\frac{2b}{N}} z^{-\frac{2}{N}(N+a-1)} \frac{\Theta_{p^N}(p^{N+b-a}q^2z^2) \Theta_{p^N}(p^N) \Theta_p(pq^2) \Theta_p(pz^2)}{\Theta_{p^N}(p^{N+b}z^2) \Theta_{p^N}(p^{N-a}q^2) \Theta_p(pq^2z^2) \Theta_p(p)}. \quad (2.12)$$

2.2 Gauge-transformed R -matrix

In order to make the comparison with our previous results easier [25, 26], one needs to introduce the following “gauge-transformed” matrix:

$$R(z, q, p) = (g^{\frac{1}{2}} \otimes g^{\frac{1}{2}}) \tilde{R}(z, q, p) (g^{-\frac{1}{2}} \otimes g^{-\frac{1}{2}}). \quad (2.13)$$

Theorem 1 *The matrix $R(z, q, p)$ satisfies the following properties:*

– *Yang-Baxter equation:*

$$R_{12}(z) R_{13}(w) R_{23}(w/z) = R_{23}(w/z) R_{13}(w) R_{12}(z), \quad (2.14)$$

– *Unitarity:*

$$R_{12}(z) R_{21}(z^{-1}) = 1, \quad (2.15)$$

– *Crossing-symmetry:*

$$R_{12}(z)^{t_2} R_{21}(z^{-1}q^{-N})^{t_2} = 1, \quad (2.16)$$

– *Antisymmetry*:

$$R_{12}(-z) = \omega(g^{-1} \otimes \mathbb{I}) R_{12}(z) (g \otimes \mathbb{I}), \quad (2.17)$$

– *Quasi-periodicity*:

$$\widehat{R}_{12}(-zp^{\frac{1}{2}}) = (g^{\frac{1}{2}}hg^{\frac{1}{2}} \otimes \mathbb{I})^{-1} \widehat{R}_{21}(z^{-1})^{-1} (g^{\frac{1}{2}}hg^{\frac{1}{2}} \otimes \mathbb{I}), \quad (2.18)$$

where

$$\widehat{R}_{12}(z) \equiv \widehat{R}_{12}(z, q, p) = \tau_N(q^{\frac{1}{2}}z^{-1}) R_{12}(z, q, p), \quad (2.19)$$

the function $\tau_N(z)$ being defined by

$$\tau_N(z) = z^{\frac{2}{N}-2} \frac{\Theta_{q^{2N}}(qz^2)}{\Theta_{q^{2N}}(qz^{-2})}. \quad (2.20)$$

The function $\tau_N(z)$ is periodic with period q^N : $\tau_N(q^N z) = \tau_N(z)$ and satisfies $\tau_N(z^{-1}) = \tau_N(z)^{-1}$.

Proof: The proof of the Yang–Baxter equation has been given in [35]. The proof of the unitarity and the crossing-symmetry is done by a direct calculation. One has to use the following two identities (the first one for the unitarity and the second one for the crossing-symmetry, see [36] for a proof of eqs. (2.21) and (2.22)):

$$\sum_{k \in \mathbb{Z}_N} S^{-i-k, i-k}(\xi, \zeta, \tau) S^{-j-k, k-j}(-\xi, \zeta, \tau) = \frac{\vartheta \left[\begin{smallmatrix} \frac{1}{2} \\ \frac{1}{2} \end{smallmatrix} \right] (\xi + \zeta, \tau) \vartheta \left[\begin{smallmatrix} \frac{1}{2} \\ \frac{1}{2} \end{smallmatrix} \right] (-\xi + \zeta, \tau)}{\vartheta \left[\begin{smallmatrix} \frac{1}{2} \\ \frac{1}{2} \end{smallmatrix} \right] (\zeta, \tau)^2} \delta_{ij}, \quad (2.21)$$

$$\sum_{k \in \mathbb{Z}_N} S^{i-k, 0}(z, \zeta, \tau) S^{j-k, 0}(-\xi - N\zeta, \zeta, \tau) = \frac{\vartheta \left[\begin{smallmatrix} \frac{1}{2} \\ \frac{1}{2} \end{smallmatrix} \right] (z, \tau) \vartheta \left[\begin{smallmatrix} \frac{1}{2} \\ \frac{1}{2} \end{smallmatrix} \right] (-\xi - N\zeta, \tau)}{\vartheta \left[\begin{smallmatrix} \frac{1}{2} \\ \frac{1}{2} \end{smallmatrix} \right] (\zeta, \tau)^2} \delta_{ij}. \quad (2.22)$$

Finally, the antisymmetry and the quasi-periodicity are explicitly checked from the expressions of the matrix elements of R . ■

Remark: The crossing-symmetry and the unitarity properties of R_{12} allow exchange of inversion and transposition for the matrix R_{12} as (the same property also holds for the matrix $\widehat{R}_{12}$):

$$\left(R_{12}(x)^{t_2} \right)^{-1} = \left(R_{12}(q^N x)^{-1} \right)^{t_2}. \quad (2.23)$$

2.3 The quantum elliptic algebra $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$

We now define the elliptic quantum algebra $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$ [1, 14] as an algebra of operators $L_{ij}(z) \equiv \sum_{n \in \mathbb{Z}} L_{ij}(n) z^n$ where $i, j \in \mathbb{Z}_N$, encapsulated into a $N \times N$ matrix

$$L(z) = \begin{pmatrix} L_{11}(z) & \cdots & L_{1N}(z) \\ \vdots & & \vdots \\ L_{N1}(z) & \cdots & L_{NN}(z) \end{pmatrix}. \quad (2.24)$$

One defines $\mathcal{A}_{q,p}(\widehat{gl}(N)_c)$ by imposing the following constraints on the $L_{ij}(z)$ (with the matrix $\widehat{R}_{12}$ given by eq. (2.19)):

$$\widehat{R}_{12}(z/w) L_1(z) L_2(w) = L_2(w) L_1(z) \widehat{R}_{12}^*(z/w), \quad (2.25)$$

where $L_1(z) \equiv L(z) \otimes \mathbb{I}$, $L_2(z) \equiv \mathbb{I} \otimes L(z)$ and $\widehat{R}_{12}^*$ is defined by $\widehat{R}_{12}^*(z, q, p) \equiv \widehat{R}_{12}(z, q, p^* = pq^{-2c})$. This definition is the most immediate generalization to N of the definition adopted in [1] for $N = 2$. The matrix $\widehat{R}_{12}^*$ obeys also the unitarity, crossing-symmetry, antisymmetry and quasi-periodicity conditions of Theorem 1 (note that the quasi-periodicity condition (2.18) for $\widehat{R}_{12}^*$ has to be understood with the modified elliptic nome p^*).

The q -determinant $q\text{-det } L(z)$ given by

$$q\text{-det } L(z) \equiv \sum_{\sigma \in \mathfrak{S}_N} \varepsilon(\sigma) \prod_{i=1}^N L_{i,\sigma(i)}(zq^{i-N-1}) \quad (2.26)$$

($\varepsilon(\sigma)$ being the signature of the permutation σ) is in the center of $\mathcal{A}_{q,p}(\widehat{gl}(N)_c)$. It can be “factored out”, and set to the value $q^{\frac{c}{2}}$ so as to get

$$\mathcal{A}_{q,p}(\widehat{sl}(N)_c) = \mathcal{A}_{q,p}(\widehat{gl}(N)_c) / \langle q\text{-det } L - q^{\frac{c}{2}} \rangle. \quad (2.27)$$

It is useful to introduce the following two matrices:

$$L^+(z) \equiv L(q^{\frac{c}{2}}z), \quad (2.28a)$$

$$L^-(z) \equiv (g^{\frac{1}{2}}hg^{\frac{1}{2}}) L(-p^{\frac{1}{2}}z) (g^{\frac{1}{2}}hg^{\frac{1}{2}})^{-1}. \quad (2.28b)$$

They obey coupled exchange relations following from (2.25) and periodicity/unitarity properties of the matrices $\widehat{R}_{12}$ and $\widehat{R}_{12}^*$:

$$\widehat{R}_{12}(z/w) L_1^\pm(z) L_2^\pm(w) = L_2^\pm(w) L_1^\pm(z) \widehat{R}_{12}^*(z/w), \quad (2.29a)$$

$$\widehat{R}_{12}(q^{\frac{c}{2}}z/w) L_1^+(z) L_2^-(w) = L_2^-(w) L_1^+(z) \widehat{R}_{12}^*(q^{-\frac{c}{2}}z/w). \quad (2.29b)$$

The parameters c, p, q in our definition are related to the corresponding parameters c', p', q' of [14] by $c = 2c'$, $p = (p')^{2/N}$, $q = (q')^{1/N}$.

3 The center of $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$ at the critical level $c = -N$

3.1 Center of $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$

Theorem 2 *At the critical level $c = -N$, the operators generated by*

$$t(z) = \text{Tr}(L(z)) = \text{Tr}\left(L^+(q^{\frac{c}{2}}z)L^-(z)^{-1}\right) \quad (3.1)$$

lie in the center of the algebra $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$.

Proof: Defining $\widetilde{L}^\pm(z) \equiv (L^\pm(z)^{-1})^t$, one can derive from eqs. (2.29) further exchange relations between the operators L^+ and $\widetilde{L}^-$:

$$\left(\widehat{R}_{12}(z/w)^{t_2}\right)^{-1} L_1^\pm(z) \widetilde{L}_2^\pm(w) = \widetilde{L}_2^\pm(w) L_1^\pm(z) \left(\widehat{R}_{12}^*(z/w)^{t_2}\right)^{-1}, \quad (3.2a)$$

$$\left(\widehat{R}_{12}(q^{\frac{c}{2}}z/w)^{t_2}\right)^{-1} L_1^+(z) \widetilde{L}_2^-(w) = \widetilde{L}_2^-(w) L_1^+(z) \left(\widehat{R}_{12}^*(q^{-\frac{c}{2}}z/w)^{t_2}\right)^{-1}. \quad (3.2b)$$

Let us now compute $\left[t(z), L^+(w)\right]$. One rewrites:

$$t(z) L^+(w) = \text{Tr}_1 \left(L_1^+(q^{\frac{c}{2}} z) \tilde{L}_1^-(z)^{t_1} \right) L_2^+(w) = \text{Tr}_1 \left(L_1^+(q^{\frac{c}{2}} z)^{t_1} \tilde{L}_1^-(z) L_2^+(w) \right) \quad (3.3)$$

since one is allowed to exchange transposition under a trace procedure. Commuting $L_2^+(w)$ through $\tilde{L}_1^-(z)$ using eq. (3.2b), one gets:

$$t(z) L_2^+(w) = \text{Tr}_1 \left(L_1^+(q^{\frac{c}{2}} z)^{t_1} (\hat{R}_{21}(q^{\frac{c}{2}} w/z)^{t_1})^{-1} L_2^+(w) \tilde{L}_1^-(z) \hat{R}_{21}^*(q^{-\frac{c}{2}} w/z)^{t_1} \right). \quad (3.4)$$

Using the unitarity property of $\hat{R}_{12}$, one has:

$$L_1^+(q^{\frac{c}{2}} z)^{t_1} (\hat{R}_{21}(q^{-\frac{c}{2}} w/z)^{-1})^{t_1} L_2^+(w) = L_2^+(w) (\hat{R}_{21}^*(q^{-\frac{c}{2}} w/z)^{-1})^{t_1} L_1^+(q^{\frac{c}{2}} z)^{t_1}. \quad (3.5)$$

Then applying eq. (2.23) to (3.5) gives:

$$L_1^+(q^{\frac{c}{2}} z)^{t_1} (\hat{R}_{21}(w q^{-\frac{c}{2}-N}/z)^{t_1})^{-1} L_2^+(w) = L_2^+(w) (\hat{R}_{21}^*(q^{-\frac{c}{2}} w/z)^{-1})^{t_1} L_1^+(q^{\frac{c}{2}} z)^{t_1}, \quad (3.6)$$

which leads at the critical value $c = -N$ to

$$L_1^+(q^{\frac{c}{2}} z)^{t_1} (\hat{R}_{21}(q^{\frac{c}{2}} w/z)^{t_1})^{-1} L_2^+(w) = L_2^+(w) (\hat{R}_{21}^*(q^{-\frac{c}{2}} w/z)^{-1})^{t_1} L_1^+(q^{\frac{c}{2}} z)^{t_1}. \quad (3.7)$$

Now, inserting (3.7) into (3.4), the first three terms in (3.4) can be rearranged and one obtains:

$$t(z) L_2^+(w) = L_2^+(w) \text{Tr}_1 \left((\hat{R}_{21}^*(q^{-\frac{c}{2}} w/z)^{-1})^{t_1} L_1^+(q^{\frac{c}{2}} z)^{t_1} \tilde{L}_1^-(z) \hat{R}_{21}^*(q^{-\frac{c}{2}} w/z)^{t_1} \right) \quad (3.8)$$

Using the fact that under a trace over the space 1 one has $\text{Tr}_1 \left(R_{21} Q_1 R'_{21} \right) = \text{Tr}_1 \left(Q_1 R'_{21}{}^{t_2} R_{21}{}^{t_2} \right)^{t_2}$, one gets

$$t(z) L_2^+(w) = L_2^+(w) \text{Tr}_1 \left(L_1^+(q^{\frac{c}{2}} z)^{t_1} \tilde{L}_1^-(z) \hat{R}_{21}^*(q^{-\frac{c}{2}} w/z)^{t_1 t_2} (\hat{R}_{21}^*(q^{-\frac{c}{2}} w/z)^{-1})^{t_1 t_2} \right)^{t_2} \quad (3.9)$$

The last two terms in the right hand side cancel each other, leaving a trivial dependence in space 2 and $\text{Tr}_1 \left(L_1^+(q^{\frac{c}{2}} z)^{t_1} \tilde{L}_1^-(z) \right) \equiv t(z)$ in space 1. This shows the commutation of $t(z)$ with $L^+(w)$ and therefore with $L^-(w) = (g^{\frac{1}{2}} h g^{\frac{1}{2}}) L^+(-p^{\frac{1}{2}} q^{-\frac{c}{2}} w) (g^{\frac{1}{2}} h g^{\frac{1}{2}})^{-1}$, hence with the full algebra $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$ at $c = -N$. $\blacksquare$

This demonstration reproduces the proof for $N = 2$ given in [25]; note that only in the $sl(N)$ crossing-symmetry relation (2.16) does N appear explicitly. The form of the operator (3.1) is identical to the form of the commuting operator derived by [37] in the case of the quantum algebra $\mathcal{U}_q(\widehat{sl}(N)_c)$. As in [25] the center of $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$ at $c = -N$ may contain other generators which we have not yet derived. However, $t(z)$ do close on their own a Poisson algebra as we are going to show.

3.2 Exchange algebra

In order to get the Poisson structure on $t(z)$, we need to compute the exchange algebra between the operators $t(z)$ and $t(w)$ when $c \neq -N$. From the definition of the element $t(z)$, one has

$$t(z)t(w) = L(z)_{i_1}^{j_1} L(w)_{i_2}^{j_2} = L^+(q^{\frac{c}{2}} z)_{i_1}^{j_1} \tilde{L}^-(z)_{i_1}^{j_1} L^+(q^{\frac{c}{2}} w)_{i_2}^{j_2} \tilde{L}^-(w)_{i_2}^{j_2}. \quad (3.10)$$

Suitable rewritings of the relations (2.29) lead to the following exchange relations between the operators L^+ and $\tilde{L}^-$:

$$\tilde{L}_1^-(z) L_2^+(w) = \left(\hat{R}_{21}(q^{\frac{c}{2}} w/z)^{t_1} \right)^{-1} L_2^+(w) \tilde{L}_1^-(z) \hat{R}_{21}^*(q^{-\frac{c}{2}} w/z)^{t_1}, \quad (3.11a)$$

$$L_1^+(z) \tilde{L}_2^-(w) = \hat{R}_{12}(q^{\frac{c}{2}} z/w)^{t_2} \tilde{L}_2^-(w) L_1^+(z) \left(\hat{R}_{12}^*(q^{-\frac{c}{2}} z/w)^{t_2} \right)^{-1}, \quad (3.11b)$$

$$\tilde{L}_1^-(z) \tilde{L}_2^-(w) = \left(\hat{R}_{12}(z/w)^{t_1 t_2} \right)^{-1} \tilde{L}_2^-(w) \tilde{L}_1^-(z) \hat{R}_{12}^*(z/w)^{t_1 t_2}. \quad (3.11c)$$

The exchange relations (2.29), (3.11) and the properties of the matrix R_{12} given in Theorem 1 then allow to move the matrices $L^+(w)$, $\tilde{L}^-(w)$ to the left of the matrices $L^+(z)$, $\tilde{L}^-(z)$. One obtains

$$t(z)t(w) = \mathcal{Y}(w/z)_{j_1 i_2}^{i_1 i_2} L(w)_{i_2}^{j_2} L(z)_{i_1}^{j_1}, \quad (3.12)$$

where $\mathcal{Y}(w/z) = T(w/z)\mathcal{M}(w/z)$ with

$$\mathcal{M}(w/z) = \left(\left(R_{21}(w/z) R_{21}(q^{c+N} w/z)^{-1} R_{12}(z/w)^{-1} \right)^{t_2} R_{12}(q^c z/w)^{t_2} \right) \quad (3.13)$$

and

$$T(w/z) = \frac{\tau_N(q^{\frac{1}{2}} z/w) \tau_N(q^{\frac{1}{2}-c} w/z)}{\tau_N(q^{\frac{1}{2}} w/z) \tau_N(q^{\frac{1}{2}-c} z/w)}. \quad (3.14)$$

3.3 Poisson structures on the center of $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$ at $c = -N$

One enounces:

Theorem 3 *The elements $t(z)$ form a closed algebra under the natural Poisson bracket on the center of $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$ given by (with $x = w/z$)*

$$\{t(z), t(w)\} = -\ln q \left(x \frac{d}{dx} \ln \tau_N(q^{\frac{1}{2}} x) - x^{-1} \frac{d}{dx^{-1}} \ln \tau_N(q^{\frac{1}{2}} x^{-1}) \right) t(z) t(w). \quad (3.15)$$

Proof: At the critical level $c = -N$, it is easy to show by direct calculation from (3.13) and (3.14) that

$$T(x)_{cr} = 1 \quad \text{and} \quad \mathcal{M}(x)_{cr} = \mathbb{I}_2 \otimes \mathbb{I}_2 \quad (3.16)$$

One recovers immediately that $t(z)t(w) = t(w)t(z)$ at the critical level $c = -N$. Hence a natural Poisson structure can be defined by

$$\{t(z), t(w)\} = \left(\frac{d\mathcal{Y}}{dc}(w/z) \right)_{j_1 j_2}^{i_1 i_2} L(w)_{i_2}^{j_2} L(z)_{i_1}^{j_1} \Big|_{cr}. \quad (3.17)$$

From eq. (3.16), one has

$$\frac{d\mathcal{Y}}{dc}(x) \Big|_{cr} = \frac{dT}{dc}(x) \Big|_{cr} \mathbb{I}_2 \otimes \mathbb{I}_2 + \frac{d\mathcal{M}}{dc}(x) \Big|_{cr}. \quad (3.18)$$

Now, one has

$$\frac{d\mathcal{M}}{dc} \Big|_{cr} = -\ln q \left\{ \left(y \frac{d}{dy} R_{21} \Big|_{y=w/z} \right)^{t_2} R_{12}(q^{-N} z/w)^{t_2} - (R_{12}(z/w)^{-1})^{t_2} \left(y \frac{d}{dy} R_{12} \Big|_{y=q^{-N} z/w} \right)^{t_2} \right\} \quad (3.19)$$

Taking now the derivative of the relation (2.23), one obtains

$$(R_{12}(x)^{t_2})^{-1} \left(y \frac{d}{dy} R_{12} \Big|_{y=x} \right)^{t_2} (R_{12}(x)^{t_2})^{-1} = \left(-y \frac{d}{dy} R_{12}^{-1} \Big|_{y=xq^N} \right)^{t_2} \quad (3.20)$$

which can be rewritten by virtue of eq. (2.23) as:

$$(R_{12}(xq^N)^{-1})^{t_2} \left(y \frac{d}{dy} R_{12} \Big|_{y=x} \right)^{t_2} (R_{12}(xq^N)^{-1})^{t_2} = \left(-y \frac{d}{dy} R_{12}^{-1} \Big|_{y=xq^N} \right)^{t_2} \quad (3.21)$$

Using then eq. (3.21) for the value $x = q^{-N}z/w$ and inserting it into eq. (3.19), one finds

$$\frac{d\mathcal{M}}{dc} \Big|_{cr} = -\ln q \left\{ \left(y \frac{d}{dy} R_{12} \Big|_{y=w/z} \right)^{t_2} + \left(y \frac{d}{dy} R_{12}^{-1} \Big|_{y=z/w} \right)^{t_2} \right\} R_{12}(q^{-N}z/w)^{t_2} \quad (3.22)$$

From the unitarity property, it follows that $\frac{d\mathcal{M}}{dc} \Big|_{cr} = 0$. Hence $\frac{d\mathcal{Y}}{dc} \Big|_{cr} = \frac{dT}{dc} \Big|_{cr} \mathbb{I}_2 \otimes \mathbb{I}_2$. This guarantees that the Poisson bracket of $t(z)$ closes on $t(z)$, a property not obvious since we have indicated that $t(z)$ may not exhaust the center of $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$.

Finally, using the q^N -periodicity property of the function τ_N and eq. (3.16), the derivative of $T(x)$ is given by:

$$\frac{dT}{dc}(x) \Big|_{cr} = \frac{d \ln T}{dc}(x) \Big|_{cr} = -\ln q \left(x \frac{d}{dx} \ln \tau_N(q^{\frac{1}{2}}x) - x^{-1} \frac{d}{dx^{-1}} \ln \tau_N(q^{\frac{1}{2}}x^{-1}) \right). \quad (3.23)$$

■

3.4 Explicit Poisson structures at $c = -N$

The structure function in eq. (3.15) is easily computed. One obtains:

$$\{t(z), t(w)\} = f(w/z) t(z) t(w), \quad (3.24)$$

where

$$\begin{aligned} f(x) = & -2 \ln q \left[\sum_{\ell \geq 0} \left(\frac{2x^2 q^{2N\ell}}{1 - x^2 q^{2N\ell}} - \frac{x^2 q^{2N\ell+2}}{1 - x^2 q^{2N\ell+2}} - \frac{x^2 q^{2N\ell-2}}{1 - x^2 q^{2N\ell-2}} \right) \right. \\ & \left. - \frac{x^2}{1 - x^2} + \frac{1}{2} \frac{x^2 q^2}{1 - x^2 q^2} + \frac{1}{2} \frac{x^2 q^{-2}}{1 - x^2 q^{-2}} - (x \leftrightarrow x^{-1}) \right]. \end{aligned} \quad (3.25)$$

We now define the Poisson structure for modes of the generating function derived from (3.24, 3.25). The modes of $t(z)$ are defined in the sense of generating functions (or formal series expansions):

$$t_n = \oint_C \frac{dz}{2\pi i z} z^{-n} t(z) \quad (3.26)$$

where C is a contour encircling the origin.

Mode expansions when the structure function $f(x)$ has an infinite set of poles at $x = q^{P(k)}$ (where $P(k)$ is integer) require a specific definition using the notion of “sectors”. This was done at the classical level in [25]. The procedure runs as follows:

The Poisson bracket between the modes is given by a double contour integral:

$$\{t_n, t_m\} = \oint_{C_1} \frac{dz}{2\pi iz} \oint_{C_2} \frac{dw}{2\pi iw} z^{-n} w^{-m} f(w/z) t(w) t(z). \quad (3.27)$$

The function $f(x)$ has here simple poles at $x = \pm q^{\pm N\ell}$ and $x = \pm q^{\pm N\ell \pm 1}$. Hence the relative position of the contours C_1 and C_2 must be specified in order to have an unambiguous result for (3.27). In addition, the antisymmetry of the Poisson brackets is only guaranteed at the mode level by an explicit symmetrization of (3.27) with respect to the position of the contours C_1 and C_2 . We shall comment on this fact when discussing the quantum mode-exchange structure. The mode Poisson bracket is thus defined as:

$$\{t_n, t_m\} = \frac{1}{2} \left(\oint_{C_1} \frac{dz}{2\pi iz} \oint_{C_2} \frac{dw}{2\pi iw} + \oint_{C_2} \frac{dz}{2\pi iz} \oint_{C_1} \frac{dw}{2\pi iw} \right) z^{-n} w^{-m} f(w/z) t(z) t(w), \quad (3.28)$$

where C_1 and C_2 are circles of radii R_1 and R_2 and one chooses $R_1 > R_2$. Explicit evaluation of (3.28) now requires to express $f(w/z)$ as a convergent Laurent series in the appropriate domains for $|w/z|$.

Let us define the sector (k) by $\frac{R_1}{R_2} \in]|q|^{-P(k)}, |q|^{-P(k+1)}[$, $P(k)$ being the ordered set of powers of q^{-1} where the poles of f are located (here $\{P(k), k \in \mathbb{N}\} = \{0, 1; N-1, N, N+1; \dots; N\ell-1, N\ell, N\ell+1; \dots\}$ respectively). For every sector (k) , eq. (3.28) defines a distinct mode Poisson bracket.

As in the $sl(2)$ case, one observes the difference between the analytic continuation formula (3.15), which is unique, and the formal series formula (3.28), where every k -labeled convergent Laurent series expansions for f may be taken as the formal series expansion of f . This fact is also mentioned in [27], considering the quantum problem.

Proposition 1 *In the case $k = 0$ (i.e. $\frac{R_1}{R_2} \in]1, |q|^{-1}[$), one finds*

$$\{t_n, t_m\}_{k=0} = -2 \ln q (q - q^{-1}) \sum_{r \in \mathbb{Z}} \frac{[(N-1)r]_q [r]_q}{[Nr]_q} t_{n-2r} t_{m+2r}, \quad (3.29)$$

where the q -numbers $[r]_q$ are defined as usual:

$$[r]_q \equiv \frac{q^r - q^{-r}}{q - q^{-1}}. \quad (3.30)$$

The proof is immediate.

When $k \neq 0$, one must add to (3.29) contributions arising from the poles at $q^{-P(j)}$ with $j = 1, \dots, k$.

Proposition 2 *The convergent series expansions in any sector (k) are obtained by adding to the coefficients of the convergent series at $k = 0$, coefficients obtained from the canonical formal series expansion of the distributions $\delta(q^{-P(j)}w/z) + \delta(-q^{-P(j)}w/z) - \delta(q^{P(j)}w/z) - \delta(-q^{P(j)}w/z)$ for $j = 1, \dots, k$, where $\delta(x) \equiv \sum_{n \in \mathbb{Z}} x^n$ for $x \in \mathbb{C}$.*

Proof: Moving from a sector (k) to a sector $(k+1)$ requires to rewrite the only term from (3.25) whose series expansion becomes divergent, namely $\frac{x^2 q^{2P(k)}}{1 - x^2 q^{2P(k)}}$, by a convergent series expansion for $|x| > |q|^{-P(k)}$. This substitutes the series $-\sum_{r \geq 0} x^{-2r} q^{-2P(k)r}$, convergent for $|x| > |q|^{-P(k)}$, to the series $\sum_{r > 0} x^{2r} q^{2P(k)r}$, convergent when $|x| < |q|^{-P(k)}$. The overall result, in the full series expansion, is to “add” the difference (order by order in x^{2r}), namely $-\sum_{r \in \mathbb{Z}} x^{2r} q^{2P(k)r} = -\delta(x^2 q^{2P(k)})$. A similar reasoning generates the term $\delta(x^2 q^{-2P(k)})$ from the x^{-1} terms in (3.25). ■

Beware that the terms obtained at $|x| = |q|^{\pm N\ell}$ get an overall 2 factor while the terms at $|x| = |q|^{\pm N\ell \pm 1}$ get an overall -1 factor.

Specification of a Poisson structure in the context of a multiple-singularities structure function therefore does require going to an explicit mode expansion.

3.5 Realization of the higher spin generators

To realize deformed $\mathcal{W}_N$ Poisson structures, we need to introduce generating functions for the higher spin objects. Having at our disposal only one commuting generating function $t(z)$, we are led by comparison with [21] to define shifted products, although with the same generator. Notice that such ordered shifted products were used a long time ago, to construct trigonometric and elliptic R -matrices from rational ones by the so-called “mean procedure” [38, 39]. We define accordingly:

$$s_i(z) = \prod_{u=-(i-1)/2}^{(i-1)/2} t(q^u z) \quad (3.31)$$

where $i = 1, \dots, N-1$.

The generators $s_1(z), \dots, s_{N-1}(z)$ then close a Poisson algebra with the following Poisson brackets $(i, j = 1, \dots, N-1)$:

$$\{s_i(z), s_j(w)\} = \sum_{u=-(i-1)/2}^{(i-1)/2} \sum_{v=-(j-1)/2}^{(j-1)/2} f\left(q^{v-u} \frac{w}{z}\right) s_i(z) s_j(w). \quad (3.32)$$

Although the generating functions (3.31) are here all constructed from one single object $t(z)$, we shall see that the Poisson structure deduced from (3.32) do not reflect this dependence and give rise indeed to genuine $\mathcal{W}_N$ -type structures, in particular recovering the q - $\mathcal{W}_N$ algebra in [21], as a consequence of the sector structure of the q - $\mathcal{W}_N$ algebra in terms of modes. Notice also from (3.32) that the limitation of the index i to values smaller than N is justified by the fact that a product of N functions $t(q^u z)$ in (3.31) in fact Poisson commute with all $s_i(w)$ owing to:

$$\sum_{u=1}^N f(q^u x) = 0, \quad \forall x \in \mathbb{C} \quad (3.33)$$

We now study the mode expansion of (3.32). The singularities of the structure function lie at $x = \pm q^{\pm N\ell} q^u$ and $\pm q^{\pm N\ell \pm 1} q^u$ with ℓ a positive integer and u an integer (resp. a half-integer) from $1 - \frac{1}{2}(i+j)$ to $\frac{1}{2}(i+j) - 1$ for $(i+j)$ even (resp. $(i+j)$ odd). They fall into sets of $(i+j+1)$ poles symmetrically arranged around $q^{\pm N\ell}$, separated by one power of q . This setting defines a labeling of sectors for the Poisson brackets as follows: for fixed i and j , we define a sector (k) by

$\frac{R_1}{R_2} \in]|q|^{-P_{ij}(k)}, |q|^{-P_{ij}(k+1)}[$, $P_{ij}(k)$ being the ordered set of powers of q^{-1} , positive and negative, where the poles of the structure function of (3.32) are located, such that $P_{ij}(0) = 0$. The rules which define the sectors in which the Poisson bracket of the modes of a given couple (s_i, s_j) is computed, are the following:

- Poisson brackets between modes of the same field $\{s_i(z), s_i(w)\}$ are required by antisymmetry to be computed on contours C_1 and C_2 symmetrized as in formula (3.28). Hence they are labeled by positive numbers (k) only, corresponding to the choice $\frac{R_1}{R_2} \in]|q|^{-P_{ii}(k)}, |q|^{-P_{ii}(k+1)}[$.
- Poisson brackets between modes of different fields $\{s_i(z), s_j(w)\}$ can be computed on a single set of contours C_1 and C_2 such that $\frac{R_1}{R_2} \in]|q|^{P_{ij}(k)}, |q|^{P_{ij}(k+1)}[$ where $P_{ij}(k)$ may be positive or negative. Symmetrization over C_1 and C_2 is not required. Antisymmetry of the Poisson bracket is imposed by computing $\{s_i, s_j\}_{(k)}$ for $i < j$ and *setting* $\{s_j, s_i\}_{(k)} \equiv -\{s_i, s_j\}_{(k)}$. Hence these Poisson brackets are labeled by positive and negative numbers, one for each couple (i, j) .
- The choice of sectors $k(i, j)$ on which Poisson brackets of different couples are computed is arbitrary. In fact, quadratic Poisson bracket structures obey the Jacobi identity as soon as they are antisymmetric, hence any antisymmetric Poisson structure is consistent.

To summarize, a complete Poisson structure for $\{s_i, s_j\}$ is characterized by the choice of $N - 1$ positive integer labels and $\frac{1}{2}(N - 1)(N - 2)$ integer labels.

Proposition 3 *When one chooses $k = 0$ for all sectors, one obtains a compact generic expression (with $n, m \in \mathbb{Z}$):*

$$\{s_i(n), s_j(m)\} = -2 \ln q (q - q^{-1}) \sum_{r \in \mathbb{Z}} \frac{[(N - \max(i, j))r]_q [\min(i, j)r]_q}{[Nr]_q} s_i(n - 2r) s_j(m + 2r). \quad (3.34)$$

The proof of this formula is given in Appendix B. This Poisson bracket structure is identical to the Poisson bracket structure obtained in [21] from a bosonization construction, excluding the extra δ -type terms in $s_{i-p} s_{j+p}$. We shall comment on the possibility of occurrence for such terms in the conclusion. This realizes, as in the $sl(2)$ case, a non-trivial connection between the $q\mathcal{W}_N$ algebra and the $sl(N)$ elliptic algebra.

Proposition 4 *Any Poisson structure in a given sector $k(i, j)$ can be obtained from (3.34) by adding to the r -dependent structure coefficient contributions from the relevant singularities of the structure function. They are given by formal power series expansions of terms $\delta(\pm q^{\pm P_{ij}(s)} w/z) s_i(z) s_j(w)$ for $s = 1, \dots, k(i, j)$.*

One should now emphasize that the Poisson structures (3.34) in any sector $k(i, j)$ are not identified to the structure which would be obtained from application by Leibniz rule to the mode expansion $\sum_{a_1 + \dots + a_i = m} \prod t_{a_i} q^{k_i}$ obtained from (3.31) by a single contour integral on a contour C_1 for z of any particular k -sector Poisson structure for the generators t_m derived in (3.29). Indeed, this structure would simply be given by the corresponding structure function $\frac{[(N - 1)r]_q [r]_q}{[Nr]_q}$ plus its δ -contributions.

By contrast, it can be seen in (3.32)-(3.34) and in the derivation (Appendix B) that (forgetting for the time being the further symmetrization requirements over the double contour integral) the Poisson structure (3.34) follows in fact from application of the Leibniz rule to expansions of the form above, but where each individual Poisson bracket $\{t_n, t_m\}$ must be computed in *distinct*

relative sectors since they stem from Poisson brackets between generating functions $t(q^{u_1}z)$ and $t(q^{u_2}w)$ given by $f(q^{u_1-u_2}z/w)$ as a contribution to the structure function, where u_1 and u_2 respectively live in two intervals $]-\frac{1}{2}(i-1), \frac{1}{2}(i-1)[$ and $]-\frac{1}{2}(j-1), \frac{1}{2}(j-1)[$: hence the relative position of integration contours depend on the difference between the indices u_1 and u_2 which lives in $]-\frac{1}{2}(i+j)-1, \frac{1}{2}(i+j)-1[$ and no individual Poisson bracket $\{t_n, t_m\}$ can be factored out. This is in particular true when considering identical fields $\{s_i, s_i\}$. The symmetrization procedure required for the t_n Poisson brackets adds a further obstacle to attempts at factoring out *symmetrized* Poisson brackets for the modes t_n .

To summarize, once the Poisson bracket of composite fields $s_i(z), s_j(w)$ are computed for the modes defined by contour integrals in specified relative positions for z and w , giving (3.32)-(3.34), the nature of composite fields $s_i(z)$ as products of the initial $t(z)$ generators (3.31) is obliterated from the new mode Poisson structure thus obtained. The composite fields then assume the nature of independent objects with the Poisson structure (3.34), thereby validating the seemingly redundant definition (3.31).

Remark: Reciprocally, all supplementary terms denoted $(\delta(\pm q^u w/z) s_i(z) s_j(w))_{n,m}$ from the mode expansion in $k \neq 0$ sectors are defined in the sense of *formal series* expansions as:

$$(\delta(\pm q^u w/z) s_i(z) s_j(w))_{n,m} \equiv \sum_{p \in \mathbb{Z}} q^{up} s_i(n-p) s_j(m+p).$$

Their Poisson brackets must therefore be computed consistently by Leibniz rule applied to their mode expansion, and *not* by using $s_i(z) s_j(q^u z)$ as a generating functional. In particular, the extra terms $\delta(q^{-1}z/w) t(z) t(w)$ in the $sl(2)$ case must not be understood as a central extension although the generating function $t(z) t(zq)$ Poisson commutes with $t(w)$. Central and lower-spin terms do not occur in our derivation. We shall comment on their absence here, and their possible reconstruction, in the conclusion.

4 Quadratic algebras in $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$

Theorem 4 *In the three-dimensional parameter space generated by p, q, c , one defines a two-dimensional surface $\Sigma_{N,M}$ for any integer $M \in \mathbb{Z}$ by the set of triplets (p, q, c) connected by the relation $(-p^{\frac{1}{2}})^{NM} = q^{-c-N}$. On the surface $\Sigma_{N,M}$, the generators $t(z)$ realize an exchange algebra with the generators $L(w)$ of $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$:*

$$t(z) L(w) = F\left(M, \frac{w}{z}\right) L(w) t(z) \quad (4.1)$$

where

$$F(M, x) = q^{2M(N-1)} \prod_{k=0}^{NM-1} \frac{\Theta_{q^{2N}}(x^{-2}p^{-k}) \Theta_{q^{2N}}(x^2p^k)}{\Theta_{q^{2N}}(x^{-2}q^2p^{-k}) \Theta_{q^{2N}}(x^2q^2p^k)} \quad \text{for } M > 0, \quad (4.2a)$$

$$F(M, x) = q^{-2|M|(N-1)} \prod_{k=1}^{N|M|} \frac{\Theta_{q^{2N}}(x^{-2}q^2p^k) \Theta_{q^{2N}}(x^2q^2p^{-k})}{\Theta_{q^{2N}}(x^{-2}p^k) \Theta_{q^{2N}}(x^2p^{-k})} \quad \text{for } M < 0. \quad (4.2b)$$

Proof: The proof runs along similar lines to the commutativity proof of Theorem 2. From eqs. (3.1) to (3.4), one gets

$$t(z) L_2^+(w) = \text{Tr}_1 \left(L_1^+(q^{\frac{c}{2}}z)^{t_1} (\widehat{R}_{21}(q^{\frac{c}{2}}w/z)^{t_1})^{-1} L_2^+(w) \widetilde{L}_1^-(z) \widehat{R}_{21}^*(q^{-\frac{c}{2}}w/z)^{t_1} \right). \quad (4.3)$$

One also has, from (3.6):

$$L_1^+(q^{\frac{c}{2}}z)^{t_1} (\widehat{R}_{21}(q^{\frac{c}{2}-c-N}w/z)^{t_1})^{-1} L_2^+(w) = L_2^+(w) (\widehat{R}_{21}^*(q^{-\frac{c}{2}}w/z)^{-1})^{t_1} L_1^+(q^{\frac{c}{2}}z)^{t_1}, \quad (4.4)$$

One realizes that the only obvious condition that allow a substitution of eq. (4.4) into eq. (4.3) using the quasi-periodicity of the matrix $\widehat{R}_{12}$ is the following:

$$(-p^{\frac{1}{2}})^{NM} = q^{-c-N} \quad \text{with } M \in \mathbb{Z} \quad (4.5)$$

Actually, from the quasi-periodicity property of $\widehat{R}_{12}$, one has:

$$\widehat{R}_{21}((-p^{\frac{1}{2}})^{NM}x) = F(M, x)\widehat{R}_{21}(x) \quad (4.6)$$

where

$$F(M, x) = \begin{cases} \prod_{k=0}^{NM-1} F\left((-p^{\frac{1}{2}})^k x\right) & \text{for } M > 0, \\ \prod_{k=1}^{N|M|} F\left((-p^{\frac{1}{2}})^{-k} x\right)^{-1} & \text{for } M < 0, \end{cases} \quad (4.7)$$

the function $F(x)$ being given by

$$F(x) = \tau_N^{-1}(q^{\frac{1}{2}}x)\tau_N^{-1}(q^{\frac{1}{2}}x^{-1}) = q^{2-2/N} \frac{\Theta_{q^{2N}}(x^2)\Theta_{q^{2N}}(x^{-2})}{\Theta_{q^{2N}}(q^2x^2)\Theta_{q^{2N}}(q^2x^{-2})}. \quad (4.8)$$

Then on the two-dimensional surface defined by (4.5), the equation (4.4) becomes

$$L_1^+(q^{\frac{c}{2}}z)^{t_1} (\widehat{R}_{21}(q^{\frac{c}{2}}w/z)^{t_1})^{-1} L_2^+(w) = F\left(M, q^{\frac{c}{2}}\frac{w}{z}\right) L_2^+(w) (\widehat{R}_{21}^*(q^{-\frac{c}{2}}w/z)^{-1})^{t_1} L_1^+(q^{\frac{c}{2}}z)^{t_1}. \quad (4.9)$$

It follows that

$$t(z) L_2^+(w) = F\left(M, q^{\frac{c}{2}}\frac{w}{z}\right) L_2^+(w) \text{Tr}_1\left((\widehat{R}_{21}^*(q^{-\frac{c}{2}}w/z)^{-1})^{t_1} L_1^+(q^{\frac{c}{2}}z)^{t_1} (L_1^-(z)^{-1})^{t_1} \widehat{R}_{21}^*(q^{-\frac{c}{2}}w/z)^{t_1}\right). \quad (4.10)$$

The two R matrices cancel due to the relation $\text{Tr}_1(R_{21}Q_1R'_{21}) = \text{Tr}_1(Q_1R'_{21}{}^{t_2}R_{21}{}^{t_2})^{t_2}$. Hence recalling that $L^+(w) = L(q^{\frac{c}{2}}w)$, one gets the desired result. Finally, one needs to compute the factor $F(M, x)$. Using eq. (4.8), one obtains the expressions of Theorem 4. ■

Theorem 5 *On the surface $\Sigma_{N,M}$, $t(z)$ closes a quadratic subalgebra:*

$$t(z)t(w) = \mathcal{Y}_{N,p,q,M}\left(\frac{w}{z}\right) t(w)t(z), \quad (4.11)$$

where

$$\mathcal{Y}_{N,p,q,M}(x) = \begin{cases} \prod_{k=1}^{NM} \frac{\Theta_{q^{2N}}^2(x^2p^{-k}) \Theta_{q^{2N}}(x^2q^2p^k) \Theta_{q^{2N}}(x^2q^{-2}p^k)}{\Theta_{q^{2N}}^2(x^2p^k) \Theta_{q^{2N}}(x^2q^2p^{-k}) \Theta_{q^{2N}}(x^2q^{-2}p^{-k})} & \text{for } M > 0, \\ \prod_{k=1}^{N|M|-1} \frac{\Theta_{q^{2N}}^2(x^2p^{-k}) \Theta_{q^{2N}}(x^2q^2p^k) \Theta_{q^{2N}}(x^2q^{-2}p^k)}{\Theta_{q^{2N}}^2(x^2p^k) \Theta_{q^{2N}}(x^2q^2p^{-k}) \Theta_{q^{2N}}(x^2q^{-2}p^{-k})} & \text{for } M < 0. \end{cases} \quad (4.12)$$

Proof: From Theorem 4, one has

$$t(z) L^+(w) = F\left(M, q^{\frac{c}{2}} \frac{w}{z}\right) L^+(w) t(z), \quad (4.13a)$$

$$t(z) (L^-(w))^{-1} = F^{-1}\left(M, -p^{\frac{1}{2}} \frac{w}{z}\right) (L^-(w))^{-1} t(z), \quad (4.13b)$$

Hence, the definition (3.1) of $t(z)$ immediately implies:

$$t(z) t(w) = \frac{F\left(M, q^{\frac{c}{2}} \frac{w}{z}\right)}{F\left(M, -p^{\frac{1}{2}} \frac{w}{z}\right)} t(w) t(z). \quad (4.14)$$

The explicit expression (4.2) for $F(M, x)$ gives the result as stated above. $\blacksquare$

We shall discuss the mode expansion of (4.11) in Sect. 6.

5 Poisson structures associated to commuting subalgebras in $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$

Theorem 6 *On the surface $\Sigma(N, M)$, when $p = q^{N^h}$ with $h \in \mathbb{Z} \setminus \{0\}$, the function $\mathcal{Y}_{N,p,q,M}$ is equal to 1. Hence $t(z)$ realizes an Abelian subalgebra in $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$.*

Proof: Theorem 6 is easily proved using the explicit expression for $F(M, x)$ and the periodicity properties of the $\Theta_{q^{2N}}$ functions. $\blacksquare$

Remark: Except in the case $N = 2$ (see [26]), no value of h allows here for $t(z)$ to be an element of a possibly extended center of $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$.

The result of Theorem 6 nevertheless allows us to define Poisson structures on the corresponding Abelian algebras. They are obtained as limits of the exchange algebra (4.11) when $p = q^{N^h}$ with $h \in \mathbb{Z} \setminus \{0\}$.

Conversely it follows that (4.14) realizes a natural quantization of the Poisson structures obtained by this limit, since it realizes an “intermediate” closed exchange algebra, contrary to the situation at $c = -N$ where $t(z)$ immediately lies in the center. This stands in contrast with the construction in [21] where the quantized $q\mathcal{W}_N$ algebras must be reconstructed by an independent quantization of the deformed classical bosons in the Cartan algebra. We see here the key role of the initial 3-parameter structure $\mathcal{A}_{q,p}(\widehat{sl}(N)_c)$ compared to the 2-parameter quantum algebra $\mathcal{U}_q(\widehat{sl}(N)_c)$ used in [21]. It allows for an intermediate 2-parameter step at $(-p^{\frac{1}{2}})^{NM} = q^{-c-N}$ where the generators $t(z)$ themselves close an exchange algebra. Hence it provides at the same time the classical q -deformed $\mathcal{W}_N$ algebra and its (q, p) -deformed quantization.

Theorem 7 *Setting $q^{N^h} = p^{1-\beta}$ for any integer $h \neq 0$, the h -labeled Poisson structure defined by:*

$$\{t(z), t(w)\}^{(h)} = \lim_{\beta \rightarrow 0} \frac{1}{\beta} \left(t(z)t(w) - t(w)t(z) \right) \quad (5.1)$$

has the following expression:

$$\{t(z), t(w)\}^{(h)} = f_h(w/z) t(z) t(w) \quad (5.2)$$

where

$$\begin{aligned}
f_h(x) &= 2Nh \ln q \left[\sum_{\ell \geq 0} E\left(\frac{NM}{2}\right) \left(E\left(\frac{NM}{2}\right) + 1\right) \left(\frac{2x^2 q^{2N\ell}}{1 - x^2 q^{2N\ell}} - \frac{x^2 q^{2N\ell+2}}{1 - x^2 q^{2N\ell+2}} - \frac{x^2 q^{2N\ell-2}}{1 - x^2 q^{2N\ell-2}} \right) \right. \\
&\quad + E\left(\frac{NM+1}{2}\right)^2 \left(\frac{2x^2 q^{2N\ell+N}}{1 - x^2 q^{2N\ell+N}} - \frac{x^2 q^{2N\ell+N+2}}{1 - x^2 q^{2N\ell+N+2}} - \frac{x^2 q^{2N\ell+N-2}}{1 - x^2 q^{2N\ell+N-2}} \right) \\
&\quad \left. - \frac{1}{2} E\left(\frac{NM}{2}\right) \left(E\left(\frac{NM}{2}\right) + 1\right) \left(\frac{2x^2}{1 - x^2} - \frac{x^2 q^2}{1 - x^2 q^2} - \frac{x^2 q^{-2}}{1 - x^2 q^{-2}} \right) - (x \leftrightarrow x^{-1}) \right] \quad \text{for } h \text{ odd}, \\
\end{aligned} \tag{5.3a}$$

$$\begin{aligned}
&= N^2 M(NM + 1) h \ln q \left[\sum_{\ell \geq 0} \left(\frac{2x^2 q^{2N\ell}}{1 - x^2 q^{2N\ell}} - \frac{x^2 q^{2N\ell+2}}{1 - x^2 q^{2N\ell+2}} - \frac{x^2 q^{2N\ell-2}}{1 - x^2 q^{2N\ell-2}} \right) \right. \\
&\quad \left. - \frac{1}{2} \left(\frac{2x^2}{1 - x^2} - \frac{x^2 q^2}{1 - x^2 q^2} - \frac{x^2 q^{-2}}{1 - x^2 q^{-2}} \right) - (x \leftrightarrow x^{-1}) \right] \quad \text{for } h \text{ even}. \\
\end{aligned} \tag{5.3b}$$

Here the notation $E(n)$ means the integer part of the number n .

Proof: direct calculation. ■

The factors $2Nh$ for h odd and $N^2 Mh(NM + 1)$ for h even are inessential and can be reabsorbed into the definition of the classical limit as $\beta \rightarrow -Nh\beta$ for h odd and $\beta \rightarrow -\frac{1}{2}N^2 M(NM + 1)h\beta$ for h even. Provided this redefinition is done, the formula (5.3b) in the case h even coincides exactly with the Poisson structure (3.25) of the center of $\mathcal{A}_{q,p}(\widehat{sl}(N)_{-N})$. The formula (5.3a) in the contrary in the case h odd gives rise to a new Poisson structure.

As before, the equations (5.3) lead to infinite families of Poisson brackets for the modes of $t(z)$ defined by (3.26). For h even, the labeling is identical to the $c = -N$ case. For h odd, the singularities lie at $\pm q^{\pm N\ell}$, $\pm q^{\pm N\ell \pm 1}$ and $\pm q^{\pm N\ell \pm \frac{1}{2}N}$, $\pm q^{\pm N\ell \pm \frac{1}{2}N \pm 1}$, giving rise to extra triplets of poles lying halfway between the initial ones.

Proposition 5 *One gets for the Poisson bracket in the $k = 0$ sector:*

$$\begin{aligned}
\{t_n, t_m\}^{(h)} &= -2 \ln q (q - q^{-1}) \sum_{r \in \mathbb{Z}} \left(E\left(\frac{NM}{2}\right) \left(E\left(\frac{NM}{2}\right) + 1\right) \frac{[(N-1)r]_q [r]_q}{[Nr]_q} \right. \\
&\quad \left. - E\left(\frac{NM+1}{2}\right)^2 \frac{[r]_q^2}{[Nr]_q} \right) t_{n-2r} t_{m+2r} \quad \text{for } h \text{ odd}, \tag{5.4a}
\end{aligned}$$

$$= -2 \ln q (q - q^{-1}) \sum_{r \in \mathbb{Z}} \frac{[(N-1)r]_q [r]_q}{[Nr]_q} t_{n-2r} t_{m+2r} \quad \text{for } h \text{ even}. \tag{5.4b}$$

A realization of higher spin generators is again achieved by the formula (3.31) with $i = 1, \dots, N-1$. Its justification will be the same as in Sect. 3.5. The generators $s_1(z), \dots, s_{N-1}(z)$ close a Poisson algebra with the following Poisson brackets ($i, j = 1, \dots, N-1$):

$$\{s_i(z), s_j(w)\}^{(h)} = \sum_{u=-(i-1)/2}^{(i-1)/2} \sum_{v=-(j-1)/2}^{(j-1)/2} f_h\left(q^{v-u} \frac{w}{z}\right) s_i(z) s_j(w). \tag{5.5}$$

The singularity structure here is as follows. Singularities of the function $f_h(x)$ occur at $x = \pm q^{\pm N\ell+u}$, $\pm q^{\pm N\ell\pm 1+u}$ where $\ell \in \mathbb{N}$ and $u \in [1 - \frac{1}{2}(i+j), \frac{1}{2}(i+j) - 1]$ when h is even. Additional singularities occur halfway between those ones, at $\pm q^{\pm N\ell \pm \frac{1}{2}N+u}$, $\pm q^{\pm N\ell \pm \frac{1}{2}N\pm 1+u}$ when h is odd. The sector structure for the Poisson brackets of $\{s_i(z), s_j(w)\}$ is easily deduced from these results.

Easiest to compute are the Poisson brackets in the sector $k = 0$ for all couples of indices (i, j) . A simpler Poisson structure in this sector is defined by furthermore taking in *all* Poisson brackets a *symmetrized* double contour integral with $\frac{R_1}{R_2} \in]1, |q|^{-1/2}[$ (or $]1, |q|^{-1}[$ depending on the parity of N and $i+j$). (Note that the symmetrized form is actually not required here either when $i \neq j$, but it leads to nicer formulae). One gets:

$$\{s_i(n), s_j(m)\}^{(h)} = -2 \ln q (q - q^{-1}) \sum_{r \in \mathbb{Z}} \frac{[(N - \max(i, j))r]_q [\min(i, j)r]_q}{[Nr]_q} s_i(n - 2r) s_j(m + 2r) \quad \text{for } h \text{ even,} \quad (5.6a)$$

$$\begin{aligned} \{s_i(n), s_j(m)\}^{(h)} = & -2 \ln q (q - q^{-1}) \sum_{r \in \mathbb{Z}} \left(E\left(\frac{NM}{2}\right) (E\left(\frac{NM}{2}\right) + 1) \frac{[(N - \max(i, j))r]_q [\min(i, j)r]_q}{[Nr]_q} \right. \\ & \left. - E\left(\frac{NM+1}{2}\right)^2 \frac{[ir]_q [jr]_q}{[Nr]_q} \right) s_i(n - 2r) s_j(m + 2r) \\ & \text{for } h \text{ odd and } i + j \leq N, \end{aligned} \quad (5.6b)$$

$$\begin{aligned} \{s_i(n), s_j(m)\}^{(h)} = & -2 \ln q (q - q^{-1}) \sum_{r \in \mathbb{Z}} \left(E\left(\frac{NM}{2}\right) (E\left(\frac{NM}{2}\right) + 1) \frac{[(N - \max(i, j))r]_q [\min(i, j)r]_q}{[Nr]_q} \right. \\ & \left. - E\left(\frac{NM+1}{2}\right)^2 \frac{[ir]_q [jr]_q}{[Nr]_q} - E\left(\frac{NM+1}{2}\right)^2 [(N - i - j)r]_q \right) s_i(n - 2r) s_j(m + 2r) \\ & \text{for } h \text{ odd and } i + j > N. \end{aligned} \quad (5.6c)$$

The Poisson bracket (5.6a) is again identical to the “core” contribution in [21] (i.e. without lower-spin extensions). (5.6b-c) however is a completely new type of quadratic q -deformed classical $\mathcal{W}_N$ -algebra with the two extra terms. It would be interesting to know about its possible explicit constructions.

6 Quantum exchange algebra

The exchange algebras (4.11) are now understood as natural quantizations of the classical q -deformed $\mathcal{W}_N$ algebras, including the initial algebra [21]. Moreover it follows that the exchange algebra (4.11) now provides us with building blocks for new deformed quantum $\mathcal{W}_{q,p}(sl(N))$ algebras with an extra integer parameter M .

We wish to describe here an explicit formulation in terms of modes of quantum generating operators $s_i(z)$ defined as in (3.31), although with a required notion of ordering between individual $t(z)$ -generators:

$$s_i(z) = \prod_{\substack{\curvearrowright \\ \frac{i-1}{2} \geq u \geq -\frac{i-1}{2}}} t(q^u z). \quad (6.1)$$

Again justification of this definition as giving genuine q - $\mathcal{W}_N$ algebras will come from the arising of

sectors in the individual $t(z)$ - $t(w)$ exchange algebra, which eventually combine in a non-trivial way in the product formula to give rise to a new algebraic structure.

The exchange algebra from (6.1) and (4.11) takes the form:

$$s_i(z)s_j(w) = \prod_{u=-\frac{i-1}{2}}^{\frac{i-1}{2}} \prod_{v=-\frac{j-1}{2}}^{\frac{j-1}{2}} \mathcal{Y}_{N,p,q,M} \left(q^{v-u} \frac{w}{z} \right) s_j(w)s_i(z), \quad (6.2)$$

for any choice of ordering in (6.1). Furthermore it follows from (6.1) and (4.12) that $s_N(z)$ commutes with all other generators; hence we are justified in restricting i to $1, \dots, N-1$.

Once we choose an exchange function in (4.12) by choosing the integer M , the next step in the procedure consists in factorizing the exchange functions in (6.2) into a function analytic around $w/z = 0$ and a function analytic around $z/w = 0$. More precisely, and following [31], one defines a Riemann problem:

$$\mathcal{Y}_{N,p,q,M;(i,j)}(x) = \mathcal{Y}_+(x)\mathcal{Y}_-^{-1}(x^{-1}) \quad (6.3)$$

in the neighborhood of a circle C of radius R . $\mathcal{Y}_+$ and $\mathcal{Y}_-$ are respectively analytic for $|x| < R$ and $|x| > R$.

Varying the values of R with respect to the position of zeroes and poles of $\mathcal{Y}$ leads to different factorizations; moreover even with fixed R the solution of (6.3) is not unique. This leads to a large choice of acceptable factorizations. It may be possible to choose $\mathcal{Y}_+ = \mathcal{Y}_-$ in the sense of analytic continuation, for instance for $i = j = 1$, when R takes any value between $|p|^{\frac{1}{2}}$ and $|p|^{-\frac{1}{2}}$; this case is treated in detail below.

The third step is specific to our approach. It consists in promoting the exchange relation deduced from (6.2) and (6.3) to the level of an analytic extension to the full complex plane of z/w :

$$\mathcal{Y}_- \left(\frac{z}{w} \right) t(z)t(w) = \mathcal{Y}_+ \left(\frac{w}{z} \right) t(w)t(z). \quad (6.4)$$

Singularities of $\mathcal{Y}_{\pm}$ now play a crucial role as in the classical case, when we define mode expansions of (6.4). The fields $s_i(z)$ are considered as abstract generating operator-valued functionals with modes defined by contour integrals:

$$s_i(n) = \oint_C \frac{dz}{2\pi iz} z^{-n} s_i(z). \quad (6.5)$$

They obey no particular supplementary relations, which may eventually arise from explicit realizations of $t(z)$ as in [27, 31].

As a consequence, we need to introduce a similar notion of “sectors”, determined by the singularities of $\mathcal{Y}_{\pm}$, and defined as the regions in the complex plane for z/w where $\mathcal{Y}_+$ and $\mathcal{Y}_-$ are given by a particular convergent series expansion (necessarily unique in *each* domain since $\mathcal{Y}_{\pm}$ are meromorphic). A choice of relative positions for the contours C_1 for z and C_2 for w thus gives the unique formal series expansions for $\mathcal{Y}_+$ and $\mathcal{Y}_-$ to be inserted in the double contour integral, which eventually gives an exchange relation for the modes. We first illustrate this on the simplest example $i = j = 1$ (i.e. $s_i(z) = s_j(z) \equiv t(z)$).

To extract from (4.11) an exchange relation between the modes t_n (3.26) of $t(z)$, we rewrite (4.11) in the form:

$$f_{N,p,q,M} \left(\frac{z}{w} \right) t(z)t(w) = t(w)t(z) f_{N,p,q,M} \left(\frac{w}{z} \right). \quad (6.6)$$

A consistent choice for $f_{N,p,q,M}(x)$ is then:

$$f_{N,p,q,M}(x) = \prod_{k=1}^{S(M)} (1-x^2)^2 \prod_{n=0}^{\infty} \frac{(1-x^2 p^{-k} q^{2N(1+n)})^2 (1-x^2 p^k q^{2+2Nn}) (1-x^2 p^k q^{-2+2N(1+n)})}{(1-x^2 p^k q^{2Nn})^2 (1-x^2 p^{-k} q^{-2+2N(1+n)}) (1-x^2 p^{-k} q^{2+2Nn})}, \quad (6.7)$$

where $S(M) = NM$ for $M > 0$ and $S(M) = N|M| - 1$ for $M < 0$.

In the hypothesis:

$$|p| < 1, \quad |q| < 1, \quad |p|^{S(M)+1} > |q|^2, \quad (6.8)$$

$f_{N,p,q,M}(x)$ is analytic for $|x| < |p|^{-\frac{1}{2}}$. Then the choice $\mathcal{Y}_+ = \mathcal{Y}_- = f_{N,p,q,M}$ solves the Riemann problem (6.3) with $|p|^{\frac{1}{2}} < R < |p|^{-\frac{1}{2}}$.

Exchange relations for the modes of $t(z)$ are then obtained by double contour integrals. We shall not give here explicit expressions of the expansion coefficients of $f_{N,p,q,M}(x)$ but we shall describe the connections between the different sector expansions.

Theorem 8 *Let $f(x)$ be a meromorphic function of x^2 having only simple and double poles, denoted by $\pm\alpha_j^{-\frac{1}{2}}$, $j \in \mathbb{N}$. Suppose that $|\alpha_1|^{-\frac{1}{2}} > 1$ and $|\alpha_j|^{-\frac{1}{2}} < |\alpha_{j'}|^{-\frac{1}{2}}$ if $j < j'$ and denote f_l the coefficients of the Taylor expansion of $f(x)$ for $|x| < |\alpha_1|^{-\frac{1}{2}}$:*

$$f(x) = \sum_{l=0}^{\infty} f_l x^{2l}. \quad (6.9)$$

Then the relation:

$$\left(\oint_{C_1} \frac{dz}{2\pi iz} \oint_{C_2} \frac{dw}{2\pi iw} + \oint_{C_2} \frac{dz}{2\pi iz} \oint_{C_1} \frac{dw}{2\pi iw} \right) z^{-n} w^{-m} [f(z/w) t(z) t(w) - f(w/z) t(w) t(z)] = 0, \quad (6.10)$$

where C_1 and C_2 are circles of radii respectively R_1 and R_2 defines a family of exchange relations for the modes t_n (3.26) of $t(z)$, depending on R_1/R_2 .

If $R_1/R_2 \in [1, |\alpha_1|^{\pm\frac{1}{2}}[$ one has:

$$\sum_{l=0}^{\infty} f_l (t_{n-2l} t_{m+2l} - t_{m-2l} t_{n+2l}) = 0, \quad (6.11)$$

where f_l is given by (6.9), while in the regions $R_1/R_2 \in]|\alpha_{j_0}|^{\pm\frac{1}{2}}, |\alpha_{j_0+1}|^{\pm\frac{1}{2}}[$, the exchange relation becomes:

$$\sum_{l \in \mathbb{Z}} f_l^{(j_0)} (t_{n-2l} t_{m+2l} - t_{m-2l} t_{n+2l}) = 0, \quad (6.12)$$

where $f_l^{(j_0)}$ ($l \in \mathbb{Z}$ and j_0 here is an index, not a power coefficient) is obtained by adding to f_l the contributions:

$$-\frac{1}{2} \alpha_j^l [(1 - \alpha_j x^2) f(x)] \Big|_{x=\alpha_j^{-1/2}}, \quad (6.13)$$

for every simple pole $\alpha_j^{-1/2}$ such that $1 \leq j \leq j_0$, and the contributions:

$$-\frac{1}{2} (l+1) \alpha_j^l [(1 - \alpha_j x^2)^2 f(x)] \Big|_{x=\alpha_j^{-1/2}} + \frac{1}{4} \alpha_j^{l-\frac{1}{2}} \frac{d}{dx} [(1 - \alpha_j x^2)^2 f(x)] \Big|_{x=\alpha_j^{-1/2}} \quad (6.14)$$

for every double pole $\alpha_j^{-1/2}$ such that $1 \leq j \leq j_0$.

Proof: If $R_1/R_2 \in [1, |\alpha_1|^{\pm\frac{1}{2}}[$, in the integrand of (6.10) both $|w/z|$ and $|z/w|$ are smaller than $|\alpha_1|^{-\frac{1}{2}}$: therefore one can develop both $f(z/w)$ and $f(w/z)$ in Taylor series according to equation (6.9). Equation (6.11) then follows immediately.

• Suppose now $R_1/R_2 \in]|\alpha_{j_0}|^{\pm\frac{1}{2}}, |\alpha_{j_0+1}|^{\pm\frac{1}{2}}[$, where $\alpha_{j_0}^{-1/2}$ is a simple pole. Without losing in generality we can suppose $R_1/R_2 \in]|\alpha_{j_0}|^{-\frac{1}{2}}, |\alpha_{j_0+1}|^{-\frac{1}{2}}[$ and rewrite eq. (6.10) in the following way:

$$\oint_{C_1} \frac{dw}{2\pi i w} \oint_{C_2} \frac{dz}{2\pi i z} z^{-n} w^{-m} [f(z/w) t(z) t(w) - f(w/z) t(w) t(z)] - (n \leftrightarrow m) = 0. \quad (6.15)$$

In the domain of integration, $f(z/w)$ can be expanded in a convergent power series of z/w according to formula (6.9), while $f(w/z)$ admits the factorization:

$$f\left(\frac{w}{z}\right) = \prod_{j=1}^{j_0-1} \frac{1}{\left(1 - \alpha_j \frac{w^2}{z^2}\right)^{m(j)}} \frac{1}{\left(1 - \alpha_{j_0} \frac{w^2}{z^2}\right)} f^{(j_0)}\left(\frac{w}{z}\right), \quad (6.16)$$

where $m(j) = 1, 2$ is the order of the j -th pole and $f^{(j_0)}(w/z)$ is an analytic function. The first two factors in the r.h.s. of (6.16) can be expanded in convergent series of z/w :

$$\prod_{j=1}^{j_0-1} \left(1 - \alpha_j \frac{w^2}{z^2}\right)^{-m(j)} = \prod_{j=1}^{j_0-1} \sum_{s=1}^{\infty} g_s^{(j)} \alpha_j^{-s} \left(\frac{z}{w}\right)^{2s}, \quad \left(1 - \alpha_{j_0} \frac{w^2}{z^2}\right)^{-1} = - \sum_{k=1}^{\infty} \alpha_{j_0}^{-k} \left(\frac{z}{w}\right)^{2k} \quad (6.17)$$

($g_s^{(j)} = -1$ if $m(j) = 1$, $g_s^{(j)} = s - 1$ if $m(j) = 2$), while $f^{(j_0)}(w/z)$ has a convergent series expansion in w/z :

$$f^{(j_0)}\left(\frac{w}{z}\right) = \sum_{l=0}^{\infty} f_l^{(j_0)} \left(\frac{w}{z}\right)^{2l}. \quad (6.18)$$

Therefore if $R_1/R_2 \in]|\alpha_{j_0}|^{\pm\frac{1}{2}}, |\alpha_{j_0+1}|^{\pm\frac{1}{2}}[$ the exchange relation (6.10) can be written, in the sense of formal power series:

$$\begin{aligned} & \sum_{l=0}^{\infty} f_l t_{n-2l} t_{m+2l} - \oint_{C_1} \frac{dw}{2\pi i w} \oint_{C_2} \frac{dz}{2\pi i z} z^{-n} w^{-m} \prod_{j=1}^{j_0-1} \sum_{s=1}^{\infty} g_s^{(j)} \alpha_j^{-s} \left(\frac{z}{w}\right)^{2s} \\ & \times \sum_{k=1}^{\infty} -\alpha_{j_0}^{-k} \left(\frac{z}{w}\right)^{2k} \sum_{l=0}^{\infty} f_l^{(j_0)} \left(\frac{w}{z}\right)^{2l} t(w) t(z) - (n \leftrightarrow m) = 0. \end{aligned} \quad (6.19)$$

Let us concentrate on the terms:

$$- \sum_{k=1}^{\infty} \alpha_{j_0}^{-k} \left(\frac{z}{w}\right)^{2k} \sum_{l=0}^{\infty} f_l^{(j_0)} \left(\frac{w}{z}\right)^{2l}. \quad (6.20)$$

This double sum can be rewritten as a sum of two contributions:

$$- \sum_{h \in \mathbb{Z}} \sum_{l=0}^{\infty} f_l^{(j_0)} \alpha_{j_0}^{h-l} \left(\frac{w}{z}\right)^{2h} + \sum_{h=0}^{\infty} \sum_{l=0}^h f_l^{(j_0)} \alpha_{j_0}^{h-l} \left(\frac{w}{z}\right)^{2h}. \quad (6.21)$$

In the second sum of (6.21) the term:

$$\sum_{l=0}^h f_l^{(j_0)} \alpha_{j_0}^{h-l} \quad (6.22)$$

is just the coefficient $f_h^{(j_0-1)}$ of the Taylor expansion of the function $f^{(j_0-1)}(w/z)$ (analytic for $\left|\frac{w}{z}\right| < |\alpha_{j_0}|^{-\frac{1}{2}}$) defined by:

$$f\left(\frac{w}{z}\right) = \prod_{j=1}^{j_0-1} \frac{1}{\left(1 - \alpha_j \frac{w^2}{z^2}\right)^{m(j)}} f^{(j_0-1)}\left(\frac{w}{z}\right). \quad (6.23)$$

Using this result, we can reinsert equation (6.21) in (6.19). We obtain:

$$\begin{aligned} & \sum_{l=0}^{\infty} f_l t_{n-2l} t_{m+2l} - \oint_{C_1} \frac{dw}{2\pi i w} \oint_{C_2} \frac{dz}{2\pi i z} z^{-n} w^{-m} \prod_{j=1}^{j_0-1} \sum_{s=1}^{\infty} g_s^{(j)} \alpha_j^{-s} \left(\frac{z}{w}\right)^{2s} \sum_{l=0}^{\infty} f_l^{(j_0-1)} \left(\frac{w}{z}\right)^{2l} t(w) t(z) \\ & + \oint_{C_1} \frac{dw}{2\pi i w} \oint_{C_2} \frac{dz}{2\pi i z} z^{-n} w^{-m} \prod_{j=1}^{j_0-1} \sum_{s=1}^{\infty} g_s^{(j)} \alpha_j^{-s} \left(\frac{z}{w}\right)^{2s} \sum_{h \in \mathbb{Z}} \sum_{l=0}^{\infty} f_l^{(j_0)} \alpha_{j_0}^{h-l} \left(\frac{w}{z}\right)^{2h} t(w) t(z) \\ & - (n \leftrightarrow m) = 0. \end{aligned} \quad (6.24)$$

The first line gives the exchange relation for $R_1/R_2 \in]|\alpha_{j_0-1}|^{\pm\frac{1}{2}}, |\alpha_{j_0}|^{\pm\frac{1}{2}}[$. The second line is the extra contribution $(\Delta_{j_0}^1)_{n,m}$ coming from the crossing of the simple singularity at $w/z = \alpha_{j_0}^{-1/2}$. The summation over h and l gives:

$$\begin{aligned} (\Delta_{j_0}^1)_{n,m} &= \oint_{C_1} \frac{dw}{2\pi i w} \oint_{C_2} \frac{dz}{2\pi i z} z^{-n} w^{-m} \delta\left(\alpha_{j_0} \frac{w^2}{z^2}\right) \prod_{j=1}^{j_0-1} \sum_{s=1}^{\infty} g_s^{(j)} \alpha_j^{-s} \left(\frac{z}{w}\right)^{2s} f^{(j_0)}\left(\alpha_{j_0}^{-1/2}\right) t(w) t(z) \\ &- (n \leftrightarrow m). \end{aligned} \quad (6.25)$$

Owing to the properties of the δ -distribution, the series expansions in (6.25) are all convergent, since $|\alpha_{j_0}/\alpha_j| < 1$ for $j < j_0$. Reinserting their sums in (6.25), one recognizes the expression:

$$\prod_{j=1}^{j_0-1} \frac{1}{\left(1 - \frac{\alpha_j}{\alpha_{j_0}}\right)^{m(j)}} f^{(j_0)}\left(\alpha_{j_0}^{-1/2}\right), \quad (6.26)$$

which, by comparison with (6.16), is just the residue of $f(x)$ at $x = \alpha_{j_0}^{-1/2}$. Then the extra contribution is:

$$(\Delta_{j_0}^1)_{n,m} = \oint_{C_1} \frac{dw}{2\pi i w} \oint_{C_2} \frac{dz}{2\pi i z} z^{-n} w^{-m} \delta\left(\alpha_{j_0} \frac{w^2}{z^2}\right) [(1 - \alpha_{j_0} x^2) f(x)] \Big|_{x=\alpha_{j_0}^{-1/2}} t(w) t(z) - (n \leftrightarrow m), \quad (6.27)$$

or after integration, using the formal power series expansion of $\delta(x)$:

$$(\Delta_{j_0}^1)_{n,m} = \sum_{l \in \mathbb{Z}} \alpha_{j_0}^l [(1 - \alpha_{j_0} x^2) f(x)] \Big|_{x=\alpha_{j_0}^{-1/2}} (t_{m-2l} t_{n+2l} - t_{n-2l} t_{m+2l}). \quad (6.28)$$

• Similar but rather longer calculations can be performed in the case of a double pole. The extra contribution $(\Delta_{j_0}^2)_{n,m}$ is now:

$$\begin{aligned}
(\Delta_{j_0}^2)_{n,m} &= \oint_{C_1} \frac{dw}{2\pi i w} \oint_{C_2} \frac{dz}{2\pi i z} z^{-n} w^{-m} \delta' \left(\alpha_{j_0} \frac{w^2}{z^2} \right) [(1 - \alpha_{j_0} x^2)^2 f(x)] \Big|_{x=\alpha_{j_0}^{-1/2}} t(w) t(z) \\
&\quad - \frac{1}{2} \oint_{C_1} \frac{dw}{2\pi i w} \oint_{C_2} \frac{dz}{2\pi i z} z^{-n} w^{-m} \delta \left(\alpha_{j_0} \frac{w^2}{z^2} \right) \alpha_{j_0}^{-1/2} \frac{d}{dx} [(1 - \alpha_{j_0} x^2)^2 f(x)] \Big|_{x=\alpha_{j_0}^{-1/2}} t(w) t(z) \\
&\quad - (n \leftrightarrow m), \tag{6.29}
\end{aligned}$$

where:

$$\delta'(x) = \sum_{h \in \mathbb{Z}} h x^{h-1}. \tag{6.30}$$

Integrating (6.29) one has:

$$\begin{aligned}
(\Delta_{j_0}^2)_{n,m} &= \sum_{l \in \mathbb{Z}} \left((l+1) \alpha_{j_0}^l [(1 - \alpha_{j_0} x^2)^2 f(x)] \Big|_{x=\alpha_{j_0}^{-1/2}} \right. \\
&\quad \left. - \frac{1}{2} \alpha_{j_0}^{l-\frac{1}{2}} \frac{d}{dx} [(1 - \alpha_{j_0} x^2)^2 f(x)] \Big|_{x=\alpha_{j_0}^{-1/2}} \right) (t_{m-2l} t_{n+2l} - t_{n-2l} t_{m+2l}). \tag{6.31}
\end{aligned}$$

Comparing (6.31,6.28) to the first line of (6.24), one obtains that (6.13,6.14) are precisely the quantities one has to add to f_l when crossing simple and double singularities. ■

Remark 1: Theorem 8 gives the mode exchange algebra between two identical fields $t(z)$ and $t(w)$ using symmetrized integration contours, see formula (6.10). A similar result can be obtained for the mode exchange algebra between two different fields $s_i(z)$ and $s_j(w)$ but with more complicated formulae. The Riemann problem (6.3) is then generically solved by two functions $\mathcal{Y}_+ \neq \mathcal{Y}_-$; the sector structure involves poles of *both* functions; the separate contributions of the poles of these two functions to the difference between adjacent sectors are given by (6.13) and (6.14) applied to the corresponding functions. The mode expansion cannot of course be factored out as in (6.12) but takes the generic form

$$\sum_{l=0}^{\infty} \mathcal{Y}_-(l) s_i(n-2l) s_j(m+2l) - \mathcal{Y}_+(l) s_j(m-2l) s_i(n+2l) = 0. \tag{6.32}$$

Finally let us emphasize that the symmetrization of the integration contours is actually not required in the quantum case, even in the case of identical fields, but it leads to nicer formulae for the coefficients $\mathcal{Y}_{\pm}(l)$.

Remark 2: It may happen that either function $\mathcal{Y}_{\pm}$, solution of (6.3), have a multiple pole α of order $\kappa > 2$. In this case, formulae analogous to (6.13) and (6.14) hold: the supplementary contribution to the coefficient f_l is then given by a sum of κ terms, each term being proportional to the n^{th} -derivative of $(1 - \alpha x^2)^{\kappa} f(x)$ taken at $x = \alpha^{-1/2}$, where $n = 0, \dots, \kappa - 1$.

Remark 3: The condition (6.8) is incompatible with the classical limit (where $q^{Nh} = p^{1-\beta}$ with $\beta \rightarrow 0$). In order to get a classical limit in the case $Nh > 2$, it is necessary to change (6.8) in such a way that the Riemann problem (6.3) is solved by two functions $\mathcal{Y}_+ \neq \mathcal{Y}_-$ (in the sense of analytic continuations).

Remark 4: Finally, let us comment on the classical limit when using *non-symmetrized* integration contours for the quantum case. One starts with the exchange formula between two different fields $s_i(z)$ and $s_j(w)$:

$$\mathcal{Y}_-(z/w) s_i(z) s_j(w) = s_j(w) s_i(z) \mathcal{Y}_+(w/z). \quad (6.33)$$

When using non-symmetrized integration contours C_1 and C_2 , one obtains for the modes

$$\sum_{l=0}^{\infty} \mathcal{Y}_-(l) s_i(n-2l) s_j(m+2l) - \mathcal{Y}_+(l) s_j(m-2l) s_i(n+2l) = 0. \quad (6.34)$$

Let us set $\mathcal{Y}_-(x) = 1 + \beta f(x)$ and $\mathcal{Y}_+(x) = 1 + \beta g(x)$. This definition is consistent since the Riemann-Hilbert problem (6.3) reads $\mathcal{Y}_+/\mathcal{Y}_- = 1$ when $\beta = 0$. (6.34) becomes:

$$\frac{1}{\beta} [s_i(n), s_j(m)] + \sum_{l \in \mathbb{Z}} f_l s_i(n-2l) s_j(m+2l) - g_l s_j(m-2l) s_i(n+2l) = 0. \quad (6.35)$$

In the classical limit $\beta \rightarrow 0$, one gets therefore, when $i \neq j$:

$$\{s_i(n), s_j(m)\} = - \sum_{l \in \mathbb{Z}} (f_l - g_{-l}) s_i(n-2l) s_j(m+2l). \quad (6.36)$$

This gives precisely the Poisson structures computed with non-symmetrized integration contours with the structure function $f(z/w) - g(w/z)$.

When $i = j$, (6.34) can be further decoupled into two exchange relations:

$$\sum_{l=0}^{\infty} (\mathcal{Y}_-(l) + \mathcal{Y}_+(l)) (s_i(n-2l) s_i(m+2l) - s_i(m-2l) s_i(n+2l)) = 0, \quad (6.37a)$$

$$\sum_{l=0}^{\infty} (\mathcal{Y}_-(l) - \mathcal{Y}_+(l)) (s_i(n-2l) s_i(m+2l) + s_i(m-2l) s_i(n+2l)) = 0. \quad (6.37b)$$

The first equation leads in the classical case to the Poisson structure

$$\{s_i(n), s_i(m)\} = - \sum_{l \in \mathbb{Z}} (f_l - g_{-l} - f_{-l} + g_l) s_i(n-2l) s_i(m+2l), \quad (6.38)$$

while the second relation has to be interpreted as supplementary constraint equations. (6.38) gives precisely the antisymmetric Poisson structure obtained in (5.4) and (5.6) by using symmetrized contour integration.

Of course, had we started with symmetrized contours in the quantum case, we would get easily the corresponding Poisson structure in the classical limit where the mode Poisson bracket is computed with symmetrized integration contours.

7 Conclusions

We have studied here exchange relations of the form $f(z/w) t(z) h(w) = h(w) t(z) f(w/z)$, and their classical Poisson bracket limits. These relations were interpreted in terms of analytic continuations and lead us to sets of formal series expansions associated to each particular convergent expansion of $f(x)$.

What we have established here is a set of universal structures for q -deformed W_N classical algebras and $\mathcal{W}_{q,p}(sl(N))$ algebras. These structures still allow for supplementary, representation-dependent extensions of the type founded in [27] or more general!

Nothing was assumed here concerning the fields $t(z)$, $h(w)$, treated as abstract objects with *a priori* no singular behaviour. Of course, had we constructed explicit realizations of these fields, we would at one stroke in the mode expansion fix a particular sector (k) *and* add possible central, linear or generically lower-spin extensions as were constructed in [27, 31].

A closer look at these particular constructions indeed shows that they all follow from bosonic realizations of the exchange algebra [21, 27] and the extensions are generated either due to cancellations between otherwise a priori independent fields in the general quadratic expression [27], or (in a different framework, that of deformed chiral algebras) due to singularities in the operator products $t(z)h(w)$ at the poles of $f(z/w)$, resolved by using explicit bosonization formulae [31].

It must therefore be expected that extended structures, with lower-spin δ -type extensions, play the most important role in practical applications of these $\mathcal{W}_{q,p}(sl(N))$ algebras. These are also crucial to define a reasonable representation theory for these algebras.

Hence the developments from the general frame which we have established are twofold. One should look for explicit realizations (bosonizations ?) of the classical/quantum algebras, thereby also getting admissible (Jacobi or cocycle solutions) extensions. Alternatively one should also look systematically for allowed extensions of our structures by defining and solving the corresponding “cocycle” equations. We hope to report on this question in a forthcoming paper [28].

Acknowledgements

This work was supported in part by CNRS, Foundation Angelo della Riccia and EC network n. FMRX-CT96-0012. L.F. is indebted to Centre de Recherches Mathématiques of Université de Montréal, where early stage of this work was done, for its kind invitation and support. M.R. and J.A. wish to thank ENSLAPP for its kind hospitality. We also wish to thank M. Jimbo and T. Miwa for a number of precious comments and indications, and drawing our attention to Ref. [14].

Appendix A: Jacobi theta functions

Let $\mathbb{H} = \{z \in \mathbb{C} \mid \text{Im}z > 0\}$ be the upper half-plane and $\Lambda_\tau = \{\lambda_1 \tau + \lambda_2 \mid \lambda_1, \lambda_2 \in \mathbb{Z}, \tau \in \mathbb{H}\}$ the lattice with basis $(1, \tau)$ in the complex plane. One denotes the congruence ring modulo N by $\mathbb{Z}_N \equiv \mathbb{Z}/N\mathbb{Z}$ with basis $\{0, 1, \dots, N-1\}$. One sets $\omega = e^{2i\pi/N}$. Finally, for any pairs $\gamma = (\gamma_1, \gamma_2)$ and $\lambda = (\lambda_1, \lambda_2)$ of numbers, we define the (skew-symmetric) pairing $\langle \gamma, \lambda \rangle \equiv \gamma_1 \lambda_2 - \gamma_2 \lambda_1$.

One defines the Jacobi theta functions with rational characteristics $\gamma = (\gamma_1, \gamma_2) \in \frac{1}{N}\mathbb{Z} \times \frac{1}{N}\mathbb{Z}$ by:

$$\vartheta \begin{bmatrix} \gamma_1 \\ \gamma_2 \end{bmatrix} (\xi, \tau) = \sum_{m \in \mathbb{Z}} \exp \left(i\pi(m + \gamma_1)^2 \tau + 2i\pi(m + \gamma_1)(\xi + \gamma_2) \right). \quad (\text{A.1})$$

The functions $\vartheta \begin{bmatrix} \gamma_1 \\ \gamma_2 \end{bmatrix} (\xi, \tau)$ satisfy the following shift properties:

$$\vartheta \begin{bmatrix} \gamma_1 + \lambda_1 \\ \gamma_2 + \lambda_2 \end{bmatrix} (\xi, \tau) = \exp(2i\pi\gamma_1\lambda_2) \vartheta \begin{bmatrix} \gamma_1 \\ \gamma_2 \end{bmatrix} (\xi, \tau), \quad (\text{A.2a})$$

$$\vartheta \begin{bmatrix} \gamma_1 \\ \gamma_2 \end{bmatrix} (\xi + \lambda_1 \tau + \lambda_2, \tau) = \exp(-i\pi\lambda_1^2 \tau - 2i\pi\lambda_1\xi) \exp(2i\pi\langle \gamma, \lambda \rangle) \vartheta \begin{bmatrix} \gamma_1 \\ \gamma_2 \end{bmatrix} (\xi, \tau), \quad (\text{A.2b})$$

where $\gamma = (\gamma_1, \gamma_2) \in \frac{1}{N}\mathbb{Z} \times \frac{1}{N}\mathbb{Z}$ and $\lambda = (\lambda_1, \lambda_2) \in \mathbb{Z} \times \mathbb{Z}$.

Moreover, for arbitrary $\lambda = (\lambda_1, \lambda_2)$ (not necessarily integers), one has the following shift exchange:

$$\vartheta \begin{bmatrix} \gamma_1 \\ \gamma_2 \end{bmatrix} (\xi + \lambda_1 \tau + \lambda_2, \tau) = \exp(-i\pi\lambda_1^2 \tau - 2i\pi\lambda_1(\xi + \gamma_2 + \lambda_2)) \vartheta \begin{bmatrix} \gamma_1 + \lambda_1 \\ \gamma_2 + \lambda_2 \end{bmatrix} (\xi, \tau). \quad (\text{A.3})$$

Considering the usual Jacobi theta function:

$$\Theta_p(z) = (z; p)_\infty (pz^{-1}; p)_\infty (p; p)_\infty, \quad (\text{A.4})$$

where the infinite multiple products are defined by:

$$(z; p_1, \dots, p_m)_\infty = \prod_{n_i \geq 0} (1 - zp_1^{n_1} \dots p_m^{n_m}), \quad (\text{A.5})$$

the Jacobi theta functions with rational characteristics $(\gamma_1, \gamma_2) \in \frac{1}{N}\mathbb{Z} \times \frac{1}{N}\mathbb{Z}$ can be expressed in terms of the Θ_p function as:

$$\vartheta \begin{bmatrix} \gamma_1 \\ \gamma_2 \end{bmatrix} (\xi, \tau) = (-1)^{2\gamma_1\gamma_2} p^{\frac{1}{2}\gamma_1^2} z^{2\gamma_1} \Theta_p(-e^{2i\pi\gamma_2} p^{\gamma_1 + \frac{1}{2}} z^2), \quad (\text{A.6})$$

where $p = e^{2i\pi\tau}$ and $z = e^{i\pi\xi}$.

Appendix B: Proof of formula (3.34)

Let us start with the equation (3.32):


$$\{s_i(z), s_j(w)\} = \sum_{u=-(i-1)/2}^{(i-1)/2} \sum_{v=-(j-1)/2}^{(j-1)/2} f\left(q^{v-u} \frac{w}{z}\right) s_i(z) s_j(w). \quad (\text{B.1})$$

where the function $f(x)$ is given by (3.25).

• Let us first suppose that $i < j$. Then, (3.32) can be rewritten as:

$$\{s_i(z), s_j(w)\} = \sum_{u=1-(i+j)/2}^{(i+j)/2-1} \eta(u) f\left(q^u \frac{w}{z}\right) s_i(z) s_j(w) \equiv f_{ij}\left(\frac{w}{z}\right) s_i(z) s_j(w), \quad (\text{B.2})$$

where the function $\eta(u)$ is given by the following graph:



Therefore, one has for the mode Poisson bracket, using symmetrized integration contours (with $m, n \in \mathbb{Z}$):

$$\{s_i(n), s_j(m)\} = \frac{1}{2} \left(\oint_{C_1} \frac{dz}{2\pi i z} \oint_{C_2} \frac{dw}{2\pi i w} + \oint_{C_2} \frac{dz}{2\pi i z} \oint_{C_1} \frac{dw}{2\pi i w} \right) z^{-n} w^{-m} f_{ij}\left(\frac{w}{z}\right) s_i(z) s_j(w). \quad (\text{B.3})$$

One has

$$\begin{aligned} f_{ij}(x) = & -2 \ln q \sum_{u=1-(i+j)/2}^{(i+j)/2-1} \eta(u) \left\{ \sum_{\ell \geq 0} \left(\frac{2x^2 q^{2N\ell+2u}}{1 - x^2 q^{2N\ell+2u}} - \frac{x^2 q^{2N\ell+2u+2}}{1 - x^2 q^{2N\ell+2u+2}} - \frac{x^2 q^{2N\ell+2u-2}}{1 - x^2 q^{2N\ell+2u-2}} \right) \right. \\ & - \sum_{\ell \geq 0} \left(\frac{2x^{-2} q^{2N\ell-2u}}{1 - x^{-2} q^{2N\ell-2u}} - \frac{x^{-2} q^{2N\ell-2u+2}}{1 - x^{-2} q^{2N\ell-2u+2}} - \frac{x^{-2} q^{2N\ell-2u-2}}{1 - x^{-2} q^{2N\ell-2u-2}} \right) - \frac{x^2 q^{2u}}{1 - x^2 q^{2u}} \\ & \left. + \frac{1}{2} \frac{x^2 q^{2u+2}}{1 - x^2 q^{2u+2}} + \frac{1}{2} \frac{x^2 q^{2u-2}}{1 - x^2 q^{2u-2}} + \frac{x^{-2} q^{-2u}}{1 - x^{-2} q^{-2u}} - \frac{1}{2} \frac{x^{-2} q^{-2u+2}}{1 - x^{-2} q^{-2u+2}} - \frac{1}{2} \frac{x^{-2} q^{-2u-2}}{1 - x^{-2} q^{-2u-2}} \right\} \quad (\text{B.4}) \end{aligned}$$

In the following, one has to distinguish the cases $i + j$ odd and $i + j$ even. In the sector $k = 0$, the integration contours C_1 and C_2 of radii R_1 and R_2 correspond to the choice $\frac{R_1}{R_2} \in]|q|^{1/2}, |q|^{-1/2}[$ when $i + j$ is odd and $\frac{R_1}{R_2} \in]|q|, |q|^{-1}[$ when $i + j$ is even (recall that $|q| < 1$). Hence, in the sector $k = 0$, the singularities of the structure function lie at $x = q^{\pm u}, q^{\pm u \pm 1}$.

a) case $i + j$ odd

In that case, u takes only half-integer values. One has therefore

$$\begin{aligned} f_{ij}(x) = & -2 \ln q \left\{ - \sum_{u=1-(i+j)/2}^{(i+j)/2-1} \eta(u) \sum_{\ell \geq 1} \sum_{r \geq 1} (q^r - q^{-r})^2 (x^{2r} q^{2N\ell r + 2ur} - x^{-2r} q^{2N\ell r - 2ur}) \right. \\ & + \sum_{r \geq 1} x^{2r} \left(\sum_{u > 0} 2\eta(u) q^{2ur} - \sum_{u > 1} \eta(u) q^{2ur-2r} - \sum_{u > -1} \eta(u) q^{2ur+2r} \right) \\ & \left. - \sum_{r \geq 1} x^{-2r} \left(\sum_{u < 0} 2\eta(u) q^{-2ur} - \sum_{u < 1} \eta(u) q^{-2ur+2r} - \sum_{u < -1} \eta(u) q^{-2ur-2r} \right) \right\} \quad (\text{B.5}) \end{aligned}$$

Then resumming the series over ℓ , (B.5) can be rewritten as:

$$f_{ij}(x) = -2 \ln q \sum_{r \geq 1} \frac{[r]_q}{[Nr]_q} (q^r - q^{-r}) \left\{ x^{2r} \left(q^{Nr} \sum_{u < 0} \eta(u) q^{2ur} + q^{-Nr} \sum_{u > 0} \eta(u) q^{2ur} - \eta\left(\frac{1}{2}\right) \frac{[Nr]_q}{[r]_q} \right) \right. \\ \left. - x^{-2r} \left(q^{-Nr} \sum_{u < 0} \eta(u) q^{-2ur} + q^{Nr} \sum_{u > 0} \eta(u) q^{-2ur} - \eta\left(\frac{1}{2}\right) \frac{[Nr]_q}{[r]_q} \right) \right\} \quad (\text{B.6})$$

From the shape of the function $\eta(u)$ (see figure above) it is easy to calculate the sums occuring in eq. (B.6). One gets

$$\sum_{u > 0} \eta(u) q^{2ur} = \sum_{u < 0} \eta(u) q^{-2ur} = \frac{q^{rj}(q^{ri} - q^{-ri}) - i(q^r - q^{-r})}{(q^r - q^{-r})^2} \\ \sum_{u > 0} \eta(u) q^{-2ur} = \sum_{u < 0} \eta(u) q^{2ur} = -\frac{q^{-rj}(q^{ri} - q^{-ri}) - i(q^r - q^{-r})}{(q^r - q^{-r})^2} \quad (\text{B.7})$$

Inserting then eq. (B.7) into eq. (B.6), one obtains

$$f_{ij}(x) = -2 \ln q (q - q^{-1}) \sum_{r \in \mathbb{Z}} -x^{2r} \frac{[(N-j)r]_q [ir]_q}{[Nr]_q} \quad (\text{B.8})$$

Hence, from eqs. (B.3) and (B.8), formula (3.34) immediately follows.

b) case $i + j$ even

In that case, u takes only integer values; in particular it can take the value $u = 0$. One has

$$f_{ij}(x) = -2 \ln q \left\{ - \sum_{u=1-(i+j)/2}^{(i+j)/2-1} \eta(u) \sum_{\ell \geq 1} \sum_{r \geq 1} (q^r - q^{-r})^2 (x^{2r} q^{2N\ell r + 2ur} - x^{-2r} q^{2N\ell r - 2ur}) \right. \\ \left. + \sum_{r \geq 1} x^{2r} \left(\sum_{u > 0} 2\eta(u) q^{2ur} - \sum_{u > 1} \eta(u) q^{2ur-2r} - \sum_{u > -1} \eta(u) q^{2ur+2r} \right) \right. \\ \left. - \sum_{r \geq 1} x^{-2r} \left(\sum_{u < 0} 2\eta(u) q^{-2ur} - \sum_{u < 1} \eta(u) q^{-2ur+2r} - \sum_{u < -1} \eta(u) q^{-2ur-2r} \right) \right\} \quad (\text{B.9})$$

Although (B.9) has the same form than (B.5), one has to be careful in the derivation of this formula due to the peculiar role of the value $u = 0$ (in particular note that $i < j$ and $i + j$ even implies $\eta(0) = \eta(\pm 1)$). When resumming the series over ℓ , one obtains after some algebra:


$$f_{ij}(x) = -2 \ln q \sum_{r \geq 1} \frac{[r]_q}{[Nr]_q} (q^r - q^{-r}) \left\{ x^{2r} \left(q^{Nr} \sum_{u < 0} \eta(u) q^{2ur} + q^{-Nr} \sum_{u > 0} \eta(u) q^{2ur} - \eta(0) \frac{[(N-1)r]_q}{[r]_q} \right) \right. \\ \left. - x^{-2r} \left(q^{-Nr} \sum_{u < 0} \eta(u) q^{-2ur} + q^{Nr} \sum_{u > 0} \eta(u) q^{-2ur} - \eta(0) \frac{[(N-1)r]_q}{[r]_q} \right) \right\} \quad (\text{B.10})$$

The sums over u are now given by:

$$\sum_{u > 0} \eta(u) q^{2ur} = \sum_{u < 0} \eta(u) q^{-2ur} = \frac{q^{rj}(q^{ri} - q^{-ri}) - iq^r(q^r - q^{-r})}{(q^r - q^{-r})^2} \\ \sum_{u < 0} \eta(u) q^{2ur} = \sum_{u > 0} \eta(u) q^{-2ur} = -\frac{q^{-rj}(q^{ri} - q^{-ri}) - iq^{-r}(q^r - q^{-r})}{(q^r - q^{-r})^2} \quad (\text{B.11})$$

Inserting (B.11) into (B.10), one gets the same formula (B.8) as in the case $i + j$ odd.

- We consider now the case $i = j$. Then the trapezoidal function $\eta(u)$ degenerates into a triangle:



The formulae (B.2)–(B.4) remain valid with the condition $i = j$. Performing the series expansions in the sector $k = 0$, one obtains

$$\begin{aligned}
f_{ii}(x) = & -2 \ln q \left\{ - \sum_{u=1-i}^{i-1} \eta(u) \sum_{\ell \geq 1} \sum_{r \geq 1} (q^r - q^{-r})^2 (x^{2r} q^{2N\ell r + 2ur} - x^{-2r} q^{2N\ell r - 2ur}) \right. \\
& + \sum_{r \geq 1} x^{2r} \left(\sum_{u > 0} 2\eta(u) q^{2ur} - \sum_{u > 1} \eta(u) q^{2ur - 2r} - \sum_{u > -1} \eta(u) q^{2ur + 2r} + 1 \right) \\
& \left. - \sum_{r \geq 1} x^{-2r} \left(\sum_{u < 0} 2\eta(u) q^{-2ur} - \sum_{u < 1} \eta(u) q^{-2ur + 2r} - \sum_{u < -1} \eta(u) q^{-2ur - 2r} + 1 \right) \right\} \quad (\text{B.12})
\end{aligned}$$

Since now $\eta(0) - \eta(\pm 1) = 1$, eq. (B.12) leads again to formula (B.10). Using (B.11), which still holds with $i = j$, one gets finally

$$f_{ii}(x) = -2 \ln q (q - q^{-1}) \sum_{r \in \mathbb{Z}} -x^{2r} \frac{[(N - i)r]_q [ir]_q}{[Nr]_q} \quad (\text{B.13})$$

which achieves the proof of (3.34).

References

- [1] O. Foda, K. Iohara, M. Jimbo, R. Kedem, T. Miwa, H. Yan, *An elliptic quantum algebra for $\widehat{sl}_2$* , Lett. Math. Phys. **32** (1994) 259.
- [2] M. Jimbo, H. Konno, T. Miwa, *Massless XXZ model and degeneration of the elliptic algebra $\mathcal{A}_{q,p}(\widehat{sl}(2)_c)$* , hep-th/9610079, Proceedings of the conference “Deformation Theory, Symplectic Geometry and Applications”, Ascona, Switzerland, 1996.
- [3] S. Khoroshkin, D. Lebedev, S. Pakuliak, *Elliptic algebra $A_{q,p}(\widehat{sl}_2)$ in the scaling limit*, q-alg/9702002.
- [4] H. Konno, *Degeneration of the elliptic algebra $A_{q,p}(\widehat{sl}_2)$ and form factors in the sine-Gordon theory*, hep-th/9701034, to appear in the CRM series in mathematical physics, Springer Verlag.
- [5] M. Jimbo, T. Miwa, A. Nakayashiki, *Difference equations for correlation functions of the eight-vertex model*, J. Phys. A **26** (1993) 2199.
- [6] M. Jimbo, R. Kedem, H. Konno, T. Miwa, R. Weston, *Difference equations in spin chains with a boundary*, Nucl. Phys. B **448** (1995) 429.
- [7] H. Konno, *An elliptic algebra $\mathcal{U}_{q,p}(\widehat{sl}(2)_c)$ and the fusion RSOS model*, q-alg/9709013.
- [8] G. Andrews, R. Baxter, P.J. Forrester, *Eight-vertex SOS model and generalized Rogers-Ramanujan type identities*, J. Stat. Phys. **35** (1984) 193.
- [9] E. Date, M. Jimbo, T. Miwa, M. Okado, *Fusion of the eight-vertex SOS model*, Lett. Math. Phys. **12** (1986) 209.
- [10] J. Avan, O. Babelon, E. Billey, *The Gervais-Neveu-Felder equation and quantum Calogero-Moser systems*, Commun. Math. Phys. **178** (1996) 281.
- [11] G. Felder, Proceedings of the ICM 1994, Birkhäuser Eds. (1994), pp. 1247.
- [12] B.Y. Hou, W.L. Yang, *Dynamically twisted algebra $\mathcal{A}_{q,p;\hat{\pi}}(\widehat{sl}(2))$ as current algebra generalizing screening currents of q -deformed Virasoro algebra*, q-alg/9709024.
- [13] C. Fronsdal, Publication RIMS Kyoto University **33** (1997) 91 and Lett. Math. Phys. **40** (1997) 117.
- [14] M. Jimbo, H. Konno, S. Odake, J. Shiraishi, *Quasi-Hopf twistors for elliptic quantum groups*, q-alg/9712029.
- [15] J.L. Gervais, A. Neveu, *Novel triangle relation and absence of tachyons in Liouville string field theory*, Nucl. Phys. B **238** (1984) 125.
- [16] D. Arnaudon, E. Buffenoir, E. Ragoucy, Ph. Roche, *Universal solutions of quantum dynamics Yang-Baxter equations*, q-alg/9712037.
- [17] J. Shiraishi, H. Kubo, H. Awata, S. Odake, *A Quantum Deformation of the Virasoro algebra and the Macdonald symmetric functions*, Lett. Math. Phys. **38** (1996) 33.

- [18] I. Andric, A. Jevicki, H. Levine, *On the large N limit in symplectic matrix models*, Nucl. Phys. B **215** (1983) 307.
- [19] J.A. Minahan, A.P. Polychronakos, *Density correlation functions in Calogero–Sutherland models*, Phys. Rev. B **50** (1994) 4236.
- [20] J. Avan, A. Jevicki, *Classical integrability and higher symmetries of collective string field theory*, Phys. Lett. B **266** (1991) 35.
- [21] E. Frenkel, N. Reshetikhin, *Quantum affine algebras and deformations of the Virasoro and W -algebras*, Commun. Math. Phys. **178** (1996) 237.
- [22] S. Lukyanov, *A note on the deformed Virasoro algebra*, Phys. Lett. B **367** (1996) 121.
- [23] S. Lukyanov, Y. Pugai, *Multipoint local height probabilities in the integrable RSOS model*, Nucl. Phys. B **473** (1996) 631.
- [24] M. Jimbo, J. Shiraishi, *A coset-type construction for the deformed Virasoro algebra*, q-alg/9709037.
- [25] J. Avan, L. Frappat, M. Rossi, P. Sorba, *Poisson structures on the center of the elliptic algebra $\mathcal{A}_{q,p}(\widehat{sl}(2)_c)$* , Phys. Lett. A **235** (1997) 323.
- [26] J. Avan, L. Frappat, M. Rossi, P. Sorba, *New $\mathcal{W}_{q,p}(sl(2))$ algebras from the elliptic algebra $\mathcal{A}_{q,p}(\widehat{sl}(2)_c)$* , q-alg/9706013, to appear in Phys. Lett. A. (1998).
- [27] B. Feigin, E. Frenkel, *Quantum W algebras and elliptic algebras*, Commun. Math. Phys. **178** (1996) 653.
- [28] J. Avan, L. Frappat, M. Rossi, P. Sorba, in preparation.
- [29] H. Awata, H. Kubo, S. Odake, J. Shiraishi, *Quantum $\mathcal{W}_N$ algebras and Macdonald polynomials*, Commun. Math. Phys. **179** (1996) 401.
- [30] B. Feigin, M. Jimbo, T. Miwa, A. Odesskii, Ya. Pugai, *Algebra of screening operators for the deformed $\mathcal{W}_N$ algebra*, q-alg/9702029.
- [31] E. Frenkel, N.Yu. Reshetikhin, *Towards deformed chiral algebras*, Proceedings of the Quantum Group Symposium at the XXIth ICGTMP, H.D. Doebner and V.K. Dobrev eds., Heron Press, Sofia (1997), pp. 27.
- [32] Y. Asai, M. Jimbo, T. Miwa, Ya. Pugai, *Bosonization of vertex operators for the $A_{N-1}^{(1)}$ face model*, J. Phys. A **29** (1996) 6595.
- [33] A.A. Belavin, *Dynamical symmetry of integrable quantum systems*, Nucl. Phys. B **180** (1981) 189.
- [34] D.V. Chudnovsky, G.V. Chudnovsky, *Completely X -symmetric S -matrices corresponding to theta functions*, Phys. Lett. A **81** (1981) 105.
- [35] C.A. Tracy, *Embedded elliptic curves and the Yang-Baxter equation*, Physica D **16** (1985) 203.

- [36] M.P. Richey, C.A. Tracy, $\mathbb{Z}_N$ *Baxter model: symmetries and the Belavin parametrization*, J. Stat. Phys. **42** (1986) 311.
- [37] N.Yu. Reshetikhin, M.A. Semenov-Tian-Shansky, *Central extensions of quantum current groups*, Lett. Math. Phys. **19** (1990) 133.
- [38] A.V. Mikhalkov, JETP Lett. **30** (1979) 443.
- [39] L. D. Fadeev, N.Yu. Reshetikhin, *The Hamiltonian structures for integrable field models*, Theor. Math. Phys. **56** (1983) 323.